

Molecular Mechanisms of DNA Damage and Repair: Progress in Plants*

Narendra Tuteja^{1, **}, Mohan B. Singh,² Mithilesh K. Misra,³
Prem L. Bhalla,² and Renu Tuteja¹

Referee: Dr. Nawin C. Mishra, Professor of Genetics, University of South Carolina,
Department of Biological Sciences, Columbia, SC 29208

¹International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi — 110 067, India; ²Institute of Land and Food Resources, The University of Melbourne, Australia; ³Department of Biochemistry, University of Lucknow, Lucknow, India

* Dedicated to Prof. Jan H. J. Hoeijmakers.

** To whom correspondence should be addressed. Phone: 91-11-6181242; Fax; 91-11-6162316; e-mail: narendra@icgeb.res.in

Table of Contents

I.	INTRODUCTION	339
II.	DNA DAMAGES	340
	A. Types of DNA Damage and Damage Products	341
	B. Free Radicals Production and Antioxidant Defense Mechanisms	343
	C. Lipid Peroxidation and DNA Damage	348
	D. Ozone and DNA Damage in Plants	350
	E. DNA Damage Response	351
III.	VARIOUS DNA REPAIR MECHANISMS	351
	A. Direct Reversal	351
	B. Base Excision Repair	353
	C. Nucleotide Excision Repair	356
	D. Photoreactivation and Photolyases	362

E.	Bypass Damage Repair	364
F.	Double-Strand Break Repair	365
G.	Mismatch Repair	369
IV.	ROLE OF DNA HELICASES IN DAMAGE AND REPAIR.....	369
V.	DESICCATION RELATED DAMAGES, TOLERANCE AND REPAIR	371
A.	In Bacterial Spores	373
B.	In Seeds	374
C.	DNA Integrity in Pollens.....	375
VI.	DNA REPAIR GENES IDENTIFIED BY SEQUENCE HOMOLOGY	375
VII.	CONCLUSION AND FUTURE PROSPECTS	377

ABBREVIATIONS: 6-4 pps, 6-4 photo products; AOS, activated oxygen species; AP, apurinic/aprimidinic; APXs, ascorbate peroxidases; BER, base excision repair; CPD, cyclobutane pyrimidine dimer; CS, Cockayne syndrome; DNA-PK, DNA-dependent protein kinase; DR, direct reversal; DSB, double strand break; ERCC, excision repair cross complementation; FEN-1, flap endonuclease; GO, 7,8-dihydro-8-oxoguanine; GR, glutathione reductase; GSH, reduced glutathione; GSSH, oxidized glutathione; HDH, human DNA helicases; L, lipid radicals; LO, alkoxy radicals; LOO, lipid peroxide; LOOH, lipid hydroperoxide; MDA, malondialdehyde; MGMT, methylguanine-methyltransferase; MMR, mismatch repair; MMS, methylmethanesulfonate; NER, nucleotide excision repair; NHEJ, nonhomologous end-joining; O₂⁻, superoxide anion radical; ·OH, hydroxyl radical; PARP-1, poly(ADP-ribose) polymerase; PCNA, proliferative cell nuclear antigen; PI3K, phosphatidylinositol 3-kinase; PUFA, polyunsaturated fatty acids; RAR, repair and recombination; RFC, replication factor C; ROS, reactive oxygen species; RPA, replication protein A; RSH, thiol radicals ('sulfur' containing functional group); SASP, small acid-soluble protein; SDSA, synthesis-dependent strand annealing; SOD, superoxide dismutase; SSA, single strand annealing; SSB, single strand break repair; UV, ultraviolet; XP, Xeroderma pigmentosum.

ABSTRACT: Despite stable genomes of all living organisms, they are subject to damage by chemical and physical agents in the environment (e.g., UV and ionizing radiations, chemical mutagens, fungal and bacterial toxins, etc.) and by free radicals or alkylating agents endogenously generated in metabolism. DNA is also damaged because of errors during its replication. The DNA lesions produced by these damaging agents could be altered base, missing base, mismatch base, deletion or insertion, linked pyrimidines, strand breaks, intra- and inter-strand cross-links.

These DNA lesions could be genotoxic or cytotoxic to the cell. Plants are most affected by the UV-B radiation of sunlight, which penetrates and damages their genome by inducing oxidative damage (pyrimidine hydrates) and cross-links (both DNA protein and DNA-DNA) that are responsible for retarding the growth and development. The DNA lesions can be removed by repair, replaced by recombination, or retained, leading to genome instability or mutations or carcinogenesis or cell death. Mostly organisms respond to genome damage by activating a DNA damage response pathway that regulates cell-cycle arrest, apoptosis, and DNA repair pathways. To prevent the harmful effect of DNA damage and maintain the genome integrity, all organisms have developed various strategies to either reverse, excise, or tolerate the persistence of DNA damage products by generating a network of DNA repair mechanisms. A variety of different DNA repair pathways have been reported that include direct reversal, base excision repair, nucleotide excision repair, photoreactivation, bypass, double-strand break repair pathway, and mismatch repair pathway. The direct reversal and photoreactivation require single protein, all the rest of the repair mechanisms utilize multiple proteins to remove or repair the lesions. The base excision repair pathway eliminates single damaged base, while nucleotide excision repair excises a patch of 25- to 32-nucleotide-long oligomer, including the damage. The double-strand break repair utilizes either homologous recombination or nonhomologous endjoining. In plant the latter pathway is more error prone than in other eukaryotes, which could be an important driving force in plant genome evolution. The *Arabidopsis* genome data indicated that the DNA repair is highly conserved between plants and mammals than within the animal kingdom, perhaps reflecting common factors such as DNA methylation. This review describes all the possible mechanisms of DNA damage and repair in general and an up to date progress in plants. In addition, various types of DNA damage products, free radical production, lipid peroxidation, role of ozone, dessication damage of plant seed, DNA integrity in pollen, and the role of DNA helicases in damage and repair and the repair genes in *Arabidopsis* genome are also covered in this review.

I. INTRODUCTION

In general, the genomes of all the living organisms, including plants, are stable. Normally the DNA of genome in the cell replicates during cell division and passes all the genetic information to their progeny. It is very important for all living organisms to ensure proper functioning and propagation of their genetic information. However, due to constant exposure of the genome to various endogenous and environmental agents, the DNA gets damaged and can produce a large variety of DNA lesions. These lesions can affect the fidelity of DNA replication (Painter, 1985), and transcrip-

tion (Protic-Sabljić and Kraemer, 1985), which can create mutations in important protein coding sequences. As a result, the produced mutated protein can affect various biological processes leading to the genome instability. Proliferating cells are often presumed to be more mutable than quiescent cells because they have less time to repair DNA damage before DNA replication. If these damages occur in the germ cells, it can be heritable and will be harmful to the next generation in passing a heritable disease. Moreover, the damage in somatic cells plays an important role in the development of cancer and aging.

The DNA damage can have genotoxic and cytotoxic effects on the cell. The real

biological consequences of these damaged products usually depend on the chemical nature of the lesion. In plants if this damage are not repaired properly they can induce proliferation as well as play an important role in the aging of seeds stocks and perennial crops. This unrepaired damage can also lead to the general deterioration of cell function and cell death. This damage situation requires constant and accurate excision and the replacement of damaged nucleotide by various DNA repair pathways. In order to maintain the integrity of the genome, the prokaryotic and eukaryotic organisms are well equipped with several DNA repair mechanism pathways (Pieper et al., 1998; Britt, 1999). In general, the lesions in the actively transcribed strand are repaired more rapidly than the lesions in the nontranscribed strand (Selby and Sancar, 1993).

DNA repair is not only a fundamental cellular process for protecting cells against the damage, but it is also essential to ensure the faithful transmission of genetic information from one generation to the next. The biological consequence of defective DNA repair in human has been reported to be the cause of many diseases, such as Xeroderma pigmentosum (XP), Cockayne syndrome (CS), and Trichothiodystrophy (TTD) (Hoeijmakers, 1993a, b; Lehmann, 1998; Winkler et al., 1998; Lindahl and Wood, 1999; Tuteja and Tuteja, 2001). However, few inherited diseases are associated with altered processing of double-strand breaks such as Ataxia telangiectasia and Nijmegen breakage syndrome (Shiloh, 1997; Lindahl and Wood, 1999). All the known mechanisms of DNA repair processes have been shown to be complex and diverse and best studied in microorganisms and mammals. In humans it is of great interest because of

the role of DNA repair in mutagenesis, in carcinogenesis, and possibly in aging (Ribeiro et al., 1998). In humans, the most important self-inflicted mutagen is tobacco smoke, which is responsible for more cancer deaths. However, the progress in plant genetic engineering and the use of mutagenesis in the creation of genetic diversity for the improvement of crops have now demanded more research in DNA damage and repair. However, DNA repair in plant has not been well studied, but now it is slowly progressing. In this article we are describing the molecular basis of DNA damage and repair and the recent advances of DNA repair in plants. Further, we have also focused on endogenous factors affecting DNA damage during natural desiccation process and the dependence of desiccation tolerance mechanisms on excision repair pathways.

II. DNA DAMAGES

The Earth's atmosphere is getting polluted due to the man-made pollutants such as chlorofluorocarbons. As a result, the stratospheric ozone layer is getting depleted, which causes the increased exposure to solar ultraviolet-B (UV-B, wavelength 280 to 320 nm) on the Earth's surface. A possible increase in UV radiation by ozone depletion is now attracting attention of researchers. This increased exposure of UV-B is harmful to all living organisms, especially to the plants. Plants are constantly being challenged by UV radiation through the sunlight because of their obligatory requirement of sunlight for photosynthesis. This radiation penetrates plant tissues and damages their genome and other cellular targets such as photosystem II and

plasma membrane ATPase (Stapleton, 1992). These ambient and increased solar radiations are responsible for retarding growth and development, morphological and biochemical alteration, and biomass accumulation in the plants (Bornman, 1989; Tevini and Teramura, 1989; Bornman and Teramura, 1993; Teramura and Sullivan, 1994; Jansen et al., 1998; Vonarx et al., 1998).

UV-B light is potentially very damaging to DNA and proteins and also increases the free radical production. Plants have their own way of tackling this problem. They have developed two protective mechanisms to cope with harmful effect of UV. First, the shielding through the production of UV-absorbing compound such as flavonoids and anthocyanins and the reflection of UV radiation by epicuticular waxes and cuticular structures (Bornman et al., 1997; Stapleton et al., 1997). Second, removal and direct reversion by photo-activation of UV-induced DNA lesions that involve photolyase (Pang and Hays, 1991; Sutherland et al., 1996; Stapleton et al., 1997; Yasui and Eker, 1998; Britt, 1999). Recently, Cathie Martin's group at John Innes Centre, UK, have shown that the production of UV-protecting sunscreens in *Arabidopsis* is controlled through the transcriptional repression by *AtMyB4*. The mutant *AtMyB4* line is more tolerant of UV-B irradiation than wild type (Jin et al., 2000). However, the light-dependent pathway is the major mechanism of repair of DNA damage in plant parts exposed to sunlight induced by UV radiation. Other pathways such as base excision repair (BER), nucleotide excision repair (NER), postreplication, and recombinational repair may be more important in internal tissues where light penetration is limited (Hoeijmakers and Bootsma, 1990;

Hoeijmakers, 1993 a, b; Wood, 1997; Balajee and Bohn, 2000).

Besides UV radiation, the plant genomic DNA is vulnerable to DNA damage from a variety of endogenous reactive metabolites, particularly oxygen-free radicals. The DNA in reproductive structures, pollen, and seed is particularly susceptible to endogenous DNA-damaging agents. In the plants there are a number of developmental stages that are particularly susceptible to the DNA damage. As a part of developmental pathway, seeds and pollens undergo the process of dehydration. This loss of the water triggers several metabolic changes, leading to free radical formations, which are capable of damaging DNA either directly or through lipid peroxidation. The accumulation of DNA lesions has been shown to accompany the seed-aging process (Boubriak et al., 1997). The nature and the role of endogenous metabolites involved in plant seed DNA damage are not clear. However, it is relevant to note that a naturally occurring product of lipid peroxidation, malondialdehyde (MDA), has been shown recently to be a major endogenous DNA damaging agent in mammalian systems (Fink et al., 1997; Marnett, 1999 a, b; Lindahl and Wood, 1999).

A. Types of DNA Damage and Damage Products

The variety of the DNA damage produced by various endogenous and exogenous agents is corrected by different repair pathways. Major DNA damage induced by different DNA damaging agents are shown in Figure 1. The endogenously generated damage to DNA is




FIGURE 1. Major DNA damage induced by different DNA-damaging agents.

known as “spontaneous DNA damage” (Britt, 1999), which is produced by the reactive metabolites (e.g., oxygen-free radicals, hydroxyl radicals, superperoxide, and nitric oxide) (Demple and Harrison, 1994) and the defect in normal processes of DNA replication or recombination. The environmental DNA-damaging agents could be UV-light, ionizing radiation, chemical mutagens, and cross-linking agents (e.g., mitomycin C and cisplatin), alkylating agents, aromatic compounds, and fungal and bacterial toxins (Plooy et al., 1984; Friedberg, 1985; Doetsch, 1995). The various DNA lesions produced by these agents are mismatches, base alterations, deletion of bases, strand breaks, cyclobutane pyrimidine dimers, 6-4 photo-products, intra- and inter-strand cross-links, alkylated bases, and bulky adducts (Figure 1) (Friedberg et al., 1995).

The most common natural environmental genotoxic agent is the UV component of sunlight. The damage induced by UV radiation to the epidermal cells of a plant is as inevitable as hydrolytic damage induced by the water present in the nucleus (Britte, 1996). Besides UV and hydrolytic damage, other damage could be alkylation damage, oxidative damage and the damage, produced by ionizing radiation (Britte, 1996). The oxidative damage is a major stress to the chilling-sensitive plants (Hodges et al., 1997; Pinehro et al., 1997). Low temperatures together with high light intensities induce the production of activated oxygen species (AOS), such as superoxide ($O_2^{\cdot-}$), hydroxyl radicals ($\cdot OH$), and hydrogen peroxide (H_2O_2) (Wise and Naylor, 1987). These radicals can react with DNA, proteins, and lipid, causing damage. However, normally the superoxide dismutase (SODs), ascorbate peroxidases (APXs), and catalases play important role as a defense against these


radicals (Inze and Van-Montagu, 1995). The environmental stresses to plants, such as drought, heat shock, salt stress, temperature, etc., can generate the production of AOS and are also known to induce plant SODs (Van Camp et al., 1994).

The chemotherapeutic agent Cisplatin forms the intrastrand adduct between the adjacent purines (AG or GG) (Plooy et al., 1984). The alkylating agents such as methylmethanesulfonate (MMS) include *N*-7-alkylguanine, *O*⁶-alkyl-guanine, *N*-3-alkyladenine, and *O*⁴-alkylthymidine induce the major damage (Beranek, 1990; Friedberg et al., 1995). The *N*-3-alkyladenine, a damage product, is highly toxic because it acts as a block to DNA replication by not recognizing the DNA polymerase (Britt, 1999). Alkylating agents can also form the phosphotriesters by alkylating the phosphodiester (Vaughan et al., 1993). The ligand Bleomycin produces single- and double-strand breaks by intercalating into DNA (Urdea et al., 1988). Overall, we can say that a single DNA-damaging agent can produce lesions that are inconsequential, mutagenic (because of mispairing), or cytotoxic (because it blocks the DNA replication or transcription). The main lesions produced in DNA by hydrolysis, reactive oxygen species, and small reactive intracellular molecules such as *S*-adenosylmethionine have been reviewed by Lindahl (1993).


B. Free Radicals Production and Anti-Oxidant Defense Mechanisms

Free radicals are chemical species (molecules or atoms) possessing an unpaired electron in their outermost orbital.

Due to the presence of one or more unpaired electrons, these species are paramagnetic, which makes them highly reactive. These are conventionally represented by a superscript dot R[•] (Dormandy, 1980). Free radicals can be formed in a molecule by gaining an additional electron, for example, the reduction of molecular oxygen (O₂) to the superoxide anion radical (O₂^{•-}).




Transition metals contain unpaired electrons and so act as free radicals, with the sole exception of zinc. The most important feature of transition metals, from the free radical point of view, is their variable valency, which allows them to undergo changes in their oxidation state involving one electron; for example, Fe (iron) has two common valencies. If a solution of ferrous salt (Iron II) is left in contact with the air, it slowly oxidizes to the ferric (Iron III) state. This is a one-electron oxidation and oxygen dissolved in solution is reduced to the superoxide radical (Halliwell and Gutteridge, 1986).








(Intermediate Complexes)

Copper (Cu) has two common valencies Cu⁺ and Cu²⁺. Similarly, under oxidation conditions copper salts can receive and donate electron to superoxide radicals O₂^{•-}



The variable valency of the transition metals helps them to be effective catalysts in many oxidation reduction reactions, and they are present for this

purpose at the active sites of many enzymes catalyzing such reactions. The most important free radicals in biological systems are a derivative of oxygen. The complete reduction of O₂ by univalent pathway results in the formation of superoxide anion, hydrogen peroxide (H₂O₂), and other products as shown below (Naqvi et al., 1986; Pryor, 1986):



Hydrogen peroxide is an oxidizing agent but not especially reactive, and its main significance lies in it being a source of hydroxyl radicals. This radical is an extremely reactive oxidizing radical that reacts with most biomolecules at diffusion-controlled rates. It will not diffuse a significant distance within a cell before reacting and has an extremely short half-life but is capable of causing a great damage within a small radius of its site of production. Under normal circumstances, the major source of free radicals in cells is electron leakage from electron transport chain, such as those in mitochondria. These react with


molecular oxygen to generate superoxide ion. Other enzymes, such as flavin oxidases located in the peroxisome, can also produce superoxide or hydrogen peroxide. Autooxidation of certain compounds, including ascorbic acid (vitamin C), thiols (e.g., Glutathione, cysteine), adrenaline, and flavin co-enzymes are yet other sources of superoxide anions in the cells. These autooxidation reactions can be greatly enhanced by the transition metal ion involvement.

Various potential sources that are responsible for the production of superoxide anions are as follows:


1. Enzymes: Various enzymes are involved in catalyzing the oxidation reactions, as a result there is univalent reduction of O_2 to O_2^- ; for example, xanthine oxidase, aldehyde oxidase, dihydrocerotic dehydrogenase, flavin dehydrogenase (Fridovich, 1976), and peroxidases (Halliwell, 1990).
2. Autooxidation: a large group of compounds, including epinephrine, flavins, and ferredoxin (Misra and Fridovich, 1972), reduced forms of riboflavin and thiol compounds such as cysteine and glutathione, generate superoxide radicals by autooxidation.
3. Subcellular organelles: the mitochondrial electron transport chain, endoplasmic reticulum substrate hydroxylation, chloroplasts, and neutrophils are all capable of producing superoxide radicals.
4. The oxidation of hemoglobin in erythrocytes is a very rich source of oxygen free radicals (Misra and Fridovich, 1972).
5. Irradiation: irradiation of living tissues also produce superoxide anion and other free radicals.

Superoxide anion and hydrogen peroxide react together to generate highly reactive hydroxyl ($\cdot OH$) radical (Beuchamp and Fridovich, 1973). Peroxides also react with metal cations to produce hydroxyl radicals in the living system. Some important mechanisms responsible for generation of hydroxyl radicals are briefly underlined.

1. Harber-Weiss reaction: in the presence of copper and/or iron (metal catalyst), H_2O_2 reacts with O_2^- and forms highly reactive $\cdot OH$ (Harber and Weiss, 1934; Kehrer, 2000).



2. Fenton reaction: damage to DNA by oxygen radicals (oxyradicals) is mediated by metal ions, especially iron, as in the classic Fenton reaction, which is the major source of highly reactive $\cdot OH$ radicals (Barb et al., 1957). This reaction was first observed by its inventor H. J. H. Fenton in 1894.




Halliwell and Gutteridge (1984) discuss the significance of Fenton reaction in free radical production in a biological system. An analogue of this reaction, where hydroperoxides replace H_2O_2 , is also important in biological systems, and is

considered to be a Fenton type reaction.



However, both the Harber-Weiss and Fenton reactions produce $\cdot\text{OH}$, but the difference is that in the Harber-Weiss reaction, metal ions are used as catalyst, while in Fenton reactions the metal ions are involved in the reaction.

3. Hydroxyl radicals are also formed during the exposure of high-energy radiations such as X-rays or γ -rays to the living tissues. Most of the energy is absorbed by the cell sap having very high water content. It may result in splitting of one of the covalent bonds of water.



Generally, free radicals in biological systems are extremely reactive and unstable. Most of these radicals exist only at a low concentration and they do not move far from their site of formation. Many free radicals are formed by the reaction of a free radical with a nonradical compound, so a chain reaction is started that may be several thousand events long, for example, lipid peroxidation involving polyunsaturated fatty acids. The primary free radical produces only local effects at the site of its production but the secondary radicals derived from it and degradation products of primary reaction cause more damaging effects at sites distinct and distant from the site where the primary free radical was formed. Two free radicals after reacting with each other form a stable molecule, which explains the

eventual termination of free radical-induced chain reaction. Reactive oxygen species (ROS), such as superoxide, H_2O_2 , and the $\cdot\text{OH}$ radical, are now known to play an important role in programmed cell death in both plants and animals (Jobs, 1999; Bethke and Jones, 2001).

Free radicals are capable of reacting with almost every known molecule of biological system in their vicinity. Free radicals damage proteins, cause breakdown of DNA strands, and initiate the peroxidation of various compounds (Rosen et al., 1993). Almost all the vital components of cells are susceptible to damage by free radicals. The damage caused by free radicals to different vital macromolecules and the mechanism involved is summarized in the Table 1.

7,8-dihydro-8-oxoguanine (GO) is a common oxidative DNA lesion generated by direct modification via reactive oxygen species. GO lesions are mutagenic and can mispair with adenine during DNA replication (Yang et al., 2001). If the resulting A/GO is not repaired before the next round of DNA replication, a C/G \rightarrow A/T transversion occurs and the opportunity for repair, is lost. The A/GO is repaired via base excision repair which is initiated by the DNA repair enzyme adenine-DNA glycosylase (Yang et al., 2001).


As a defense against the deleterious actions of free radicals, cells contain several enzymes for removing free radicals and their products. These are known as the antioxidant defense system or as free radical scavengers (Sies, 1991). Antioxidant means "against oxidation". Antioxidants work to protect lipids from peroxidation by radicals. Antioxidants are effective because they give up their own electrons to free radicals and thereby break the oxidation chain reaction. Some of the important ones are superoxide

TABLE 1
Biological Roles of Oxy Free Radicals

Radical species	Damaging roles	Protectors
$O_2^{\cdot -}$	RSH damage Enzyme activation and inactivation Fenton reaction	SOD
$O_2^{\cdot -}$	Lipid peroxidation Fenton-type reaction	Vit. C and E β -carotene
$\cdot OH$	Breaking of DNA strand Lipid peroxidation Secondary radical peroxidation	RSH antioxidants DNA repair process
$RO^{\cdot -}$	Secondary radical peroxidation	RSH antioxidants
$ROOH$	Peroxidation Fenton reaction	Glutathione peroxidase
H_2O_2	Peroxidase action Fenton reaction	Catalase

dismutase (SOD), catalase, and glutathione peroxidase.


Superoxide dismutase (SOD): This enzyme directly interacts with oxy free radical to convert it to hydrogen peroxide.



Cytosolic enzyme contains copper and zinc (Cu/Zn-SOD) (Fee, 1982), whereas mitochondrial enzyme contains Mn (Mn-SOD) (Weisigner and Fridovich, 1973). In plant cells the active oxygen species (AOSs) are also known to be produced under stress conditions. Recently, in rice seedling the responses of SOD genes to environmental stresses, such as drought, salinity, and chilling, were analyzed by Kaminaka et al. (1999). They observed that the expression of abscisic acid (ABA)-inducible genes, Mn-SOD gene (*sodA1*), and one of the cytosolic Cu/Zn-SOD gene (*sodCc2*) were strongly induced under the drought and salinity stresses. While Fe-SOD gene

(*sodB*) and the other cytosolic Cu/Zn-SOD gene (*sodCc1*) were also reported to be induced by ABA. However, the mRNA level of *sodB* was decreased under drought stress, and *sdCc1* gene did not induce under drought and stress conditions. The plastidic Cu/Zn-SOD gene (*sodcp*) altered under salinity stresses in the light but not in the dark. The above results clearly showed that phytohormone and AOSs are associated with the regulation of SOD genes under environmental stresses (Kaminaka et al., 1999).


Catalase: The enzyme catalyses the transformation of less toxic hydrogen peroxide to water




It is a heme-containing enzyme present in the peroxisomes and the other cell organelles.

Glutathione peroxidase: This enzyme catalyzes the oxidation of reduced glutathione (GSH) to oxidized form (GSSG) at the expense of H_2O_2 thereby remov-

ing toxic H₂O₂. Furthermore, glutathione peroxidase also metabolizes lipid hydroperoxides to hydroxyl fatty acids



The supply of reduced glutathione for continuous removal of H₂O₂ is maintained by the glutathione reductase



In plants free radicals are known to be present *in vivo* in the young roots enriched with oxygen (Goodman et al., 1986), in the germinating seeds subjected to dehydration (Senaratna and McKersie, 1986), in ungerminated seed and pollen, and also accumulate during artificial aging (Priestley et al., 1985). As a result of the drought-induced impairment of the electron transport system, the superoxide radical is generated, which damages the membrane in droughted wheat (Halliwell, 1987; Hendry et al., 1989; Price et al., 1989). It was known that through Fenton reaction, the superoxide is reduced to the highly reactive hydroxyl radical, which causes severe peroxidative damage to lipid membranes (Price and Hendry, 1989; Price et al., 1989). The extracellular sources of activated oxygen could be air pollutants such as ozone (Mehlhorn et al., 1990; Kanofsky and Sima, 1991) and the radicals produced by neighbouring cells during the hypersensitive response (Levine et al., 1994).


Recently, Luxford et al. (1999) demonstrated conclusively that Histone H1 and other protein and amino acid hydroperoxides can give rise to free radicals, which might result in oxidative damage to associated DNA. The hydroperoxides

can decompose to oxygen- and carbon-centred radicals (detected by electron paramagnetic resonance spectroscopy) on exposure to Cu⁺ and other transition metal ions. These hydroperoxide-derived radicals react readily with pyrimidine DNA bases and nucleosides to give adduct species (i.e., protein-DNA base cross-links). Product analysis has demonstrated that radicals from histone H1-hydroperoxides and other protein and amino acid hydroperoxides can also oxidize both free 2'-deoxyguanosine and intact calf thymus DNA to give the mutagenic oxidized base 8-oxo-2'-deoxyguanosine (8-oxo dG). 8-oxo dG is considered to be a precise and sensitive biomarker of oxidative DNA damage (Liu and Wells, 1995). Bialkowski and Olinski (1999) studied 8-oxo dG levels in DNA as a measure of oxidative DNA damage in Cardamine pratensis plant as affected by different environmental parameters such as light, water accessibility, and low temperature. No significant difference in 8-oxo dG level between DNA of etiolated and light-exposed plants and between DNA of drought and control plants was found. However, cold temperature (1°C for 28 h)-treated plants showed an increase of 8-oxo dG in their DNA, indicating oxidative damage to plant DNA under low-temperature conditions.

C. Lipid Peroxidation and DNA Damage


Lipid peroxidation is a highly destructive process that causes DNA damage and alters RNA transport from nucleus to cytoplasm. It also alters the structure and function of cellular membrane (Agrawal and Kale, 2001). All

the major classes of biomolecules may be attacked by the free radicals, but lipids are probably the most susceptible. The cell membranes are rich sources of polyunsaturated fatty acids (PUFA), which are readily attacked by oxidizing radicals. The oxidative destruction of PUFAs, known as lipid peroxidation (Tappel, 1973), causes a loss of fluidity and a breakdown of the membrane's secretory functions and transmembrane ionic gradient chain reaction (Barber and Berheim, 1967). They can further oxidize PUFA molecules and initiate new chain-producing lipid hydroperoxides (LOOH) that can break down to yet more radicals and to a wide range of compounds, notably aldehydes (Esterbauer et al., 1990a). Therefore, lipid peroxidation can be used as a measure of oxidative damage.



The end products of lipid peroxidation processes are aldehydes, hydrocarbon gases, and various chemical residues (del Maestro et al., 1981a). These degradation products can diffuse away from the site of chain reaction and give rise to cell edema, influence vascular permeability, and cause inflammation and chemotaxis in animals. These products may also alter the activity of phospholipids and induce the release of arachidonic acid with the subsequent formation of prostaglandins (Hemler and Lands, 1980) and various endoperoxides (del Maestro et al., 1981a, b).

The $\text{O}_2\cdot$ can indirectly influence lipid peroxidation *via* Fenton/Harber-Weiss reaction.



The generation of $\text{O}_2\cdot$ by any source, in the presence of metal ions, particularly iron ions, can lead to the formation of $\cdot\text{OH}$ radical and in turn may initiate lipid peroxidation. However, lipid peroxidation has no intrinsic metal ion requirement. Nonetheless, iron compounds can increase the rate of propagation of lipid peroxidation dramatically by decomposing lipid hydroperoxides (LOOH) to reactive alkoxy ($\text{LO}\cdot$) or peroxy ($\text{LOO}\cdot$) radicals, which can initiate new reactions (Agrawal and Kale, 2001).

Lipid peroxidation causes the endogenous lesions by inducing the formation of exocyclic DNA base adducts (Lindahl and Wood, 1999). A major recent advancement has been the detection of number of lesions in DNA that can be attributed to metabolic sources (Fink et al., 1997). Lipid peroxidation, for example, leads to the formation of a complex variety of nucleophiles capable of reacting with proteins and DNA (reviewed by Marnett, 1999a,b). The reaction of nucleophiles with DNA leads to formation of mutagenic adducts. The lipid peroxidation product malondialdehyde (MDA) reacts with G residue in DNA to form a pyridimidopurinone called M_1G , an exocyclic adduct that can block the Watson-Crick base pairing of DNA bases and thus have the potential to block DNA replication (Marnett, 1999b). Site-directed mutagenesis studies have shown that M_1G is an efficient premutagenic lesion (Fink et al., 1997). The M_1G adduct is not very stable and can decompose to a secondary ring-

opened derivative. In addition to MDA, lipid peroxidation can also produce acrolein and crotonaldehyde, which are readily metabolized to epoxides. This resulted in the generation of exocyclic etheno modifications of DNA bases (Lindahl and Wood, 1999). Two such bases, etheno-A and etheno-C, are removed by DNA glycosylases (Hang et al., 1997; Saparbaev and Laval, 1998). These strongly suggest that the generation of such adducts occurs at high rate (Lindahl and Wood, 1999).

Several leading laboratories around the world are currently addressing the identification of endogenous DNA lesions in mammalian DNA and their role in genetic disease. As lipid peroxidation and DNA damage individually have been considered major determinants of seed viability loss; however, no linkage between lipid peroxidation product and DNA damage in seed deterioration has been contemplated so far.

D. Ozone and DNA Damage

Ozone (O_3) itself is not a free radical, but the damage caused by it is often mediated by free radical production and thereby can stimulate lipid peroxidation. In last 40 years, the tropospheric O_3 concentration has increased 2- to 5-fold (Kley et al., 1999). Exposure of animals to O_3 is known to damage lungs (Mustafa and Tiernery, 1978). However, O_3 poses a 2-fold challenge to plants: photosynthesis and growth can be impaired and it can also produce foliar lesions in sensitive species and cultivars (Kley et al., 1999). O_3 has also been regarded as a "wound stress" that causes necrosis by oxidizing and damaging plasma membranes (reviewed in Heath and Taylor,

1997). O_3 responses are known to resemble components of the hypersensitive response (HR) found in incompatible plant-pathogen interactions (Sharma and Davis, 1997; Sandermann et al., 1998). This resemblance is most likely related to the occurrence of relative oxygen species (ROS), such as superoxide anion radicals ($O_2^{\cdot -}$) and H_2O_2 in the apoplast, which trigger an oxidative burst in the affected cells (Pellinen et al., 1999; Rao and Davis, 1999). H_2O_2 has been reported to be accumulated in response to O_3 in tobacco (Schraudner et al., 1998) and birch (Pellinen et al., 1999). O_3 stress in *Arabidopsis* is reported to be responsible for cell death (Rao and Davis, 1999).

O_3 also activates ethylene and salicylate (SA) signal transduction pathways leading to downstream responses, such as antioxidant and antimicrobial defenses (Sharma and Davis, 1997; Sandermann et al., 1998). The toxicity of O_3 is through chemical reaction with ethylene, yielding toxic products that initiate a self-propagating lipid peroxidation (Mehlhorn and Wellburn, 1987). The emission of ethylene is a kind of wounding symptom in O_3 -stressed plants (Heath and Taylor, 1997). In pea carpel senescence, maize endosperm development, as well as in root aerenchyma formation, ethylene is reported to regulate the programmed cell death (Orzaez and Granell, 1997; Young et al., 1997; Drew et al., 2000). Recently, Overmyer et al. (2000) have isolated an *Arabidopsis* mutant, *rcdl* (for radical-induced cell death 1), which displayed O_3 - and $O_2^{\cdot -}$ -inducible lesion formation. These authors proposed a model for the relative contribution of the different signaling pathways to ROS-driven lesion propagation and the process of lesion containment.

E. DNA Damage Response

Most organisms respond to chromosomal insults by activating a complex damage response pathway that regulates some known responses such as cell-cycle arrest, apoptosis, and direct activation of repair networks (Zhou and Elledge, 2000). DNA damage checkpoints control the ability of cells to arrest the cell cycle in response to DNA damage. The DNA damage response pathway is a kind of signal transduction pathway consisting of sensor, transducers, and effectors. The identities of sensors are not yet known, but the DNA-break binding protein such as poly (ADP-ribose) polymerase (PARP) and DNA-dependent protein kinase (DNA-PK) are the kind of DNA damage sensors. Much is known about signal transducers, which are composed of four sets of conserved proteins with some recognizable motifs. These four classes are

1. Phospho-inositide kinase (PIK)-related proteins, which include ATM and ATM-Rad3-related (ATR) in mammals and their homologs in yeast (Mec1, Rad3, Tel1). Downstream of these proteins are two check point kinases (chk1 and chk2).
2. BRCT-repeat containing proteins, which include budding yeast Rad9 and fission yeast Crb2.
3. RFC-like proteins, for example, Rad17, RFC2.5, in mammals and Rad24, Rad17, RFC2-5 and RFC3 in yeasts.
4. PCNA-like proteins, for example, Rad1, Rad9, Hus1 in mammals and Rad17, Rad1, Ddc1, Rad9, Mec3, and Hus1 in yeasts.

The effectors of damage response could be either activation of DNA repair

pathways or cell cycle arrest or apoptosis depending on the severity and type of the damage (Zhou and Elledge, 2000).

III. VARIOUS DNA REPAIR MECHANISMS

For more than 2.5 billion years since the first appearance of living cells on Earth, they were continuously exposed to solar UV radiation and many DNA damaging agents. To protect the DNA from various damage, an intricate network of DNA repair mechanisms has evolved early in evolution (de Laat et al., 1999). Several DNA repair pathways known to date in different living cells are shown in Figure 2. Out of these, the nucleotide excision repair (NER) has been studied extensively. Although the DNA repair mechanisms have been described thoroughly in *E. coli*, *Saccharomyces cerevisiae*, rodents, and human, little is known still about these processes in plants. Any alteration in the DNA double helix, even as small as single base damage, are easily recognized. Such recognition is usually by a change in the physical structure of the DNA double helix. The repair mechanisms protect the genome by repairing modified bases, DNA adducts, crosslinks, and strand breaks. Furthermore, these repair mechanisms are also coordinated with other cellular functions, in particular gene transcription and cell cycle. The various basic repair mechanisms by which damages are eliminated from DNA are described below.

A. Direct Reversal (DR)

DR is a simple mechanism that involves a single-enzyme reaction for the




FIGURE 2. Several DNA repair pathways used by different living cells. **MGMT** = methylguanine methyltransferase; **DR** = direct reversal; **BER** = base excision repair; **NER** = nucleotide excision repair; **SSBR** = single strand break repair; **DSBR** = double-strand break repair; **MMR** = mismatch repair; **PR** = photoreactivation.

removal of certain types of DNA damage. Alkyltransferases simply extract the alkyl group from alkylated bases that is transferred to an internal cysteine residue, and thus inactivate themselves (Teo et al., 1984). The best example for DR is the correction of the miscoding alkylation lesion *O*⁶-methylguanine, which is generated endogenously in small amounts by reactive cellular catabolites (Vaughan et al., 1993). DR is carried out by a specific enzyme, called methylguanine-methyltransferase (MGMT), which removes the methyl group from the guanine residue of DNA, and transfers it to one of its own cysteine residues (Figure 2) in a rapid and error-free repair process (Moore et al., 1994). *O*⁶-methyl-guanine can pair with both C and T and thereby cause transition mutations, which are sometime corrected by mismatch repair mechanism (Lindahl and Wood, 1999).

Photolyases, on the other hand, revert UV-induced dimers in a light-dependent reaction called photo-reactivation (PR) (Sancar, 1990; Yasui and Eeker 1998; Todo, 1999), which is discussed later.

B. Base Excision Repair (BER)

The BER mechanism is known for the elimination of single damaged base residues in DNA and is considered to be an essential pathway for DNA maintenance. BER mainly removes the DNA damages that are arising spontaneously in a cell from hydrolytic events such as deamination or base loss, fragmented bases resulting from ionizing radiation, and oxidative damages or methylation of ring nitrogens by endogenous agents. BER pathways involve several steps,

which include recognition and removal of damaged base, incision, gap filling, and sealing (Britt, 1996; Lehman, 1998; Lindahl and Wood, 1999). A model of various steps in BER is shown in Figure 3. The first step in BER involves the removal of a single damaged base through the action of a specific DNA *N*-glycosylase, which hydrolytically cleaves the base-deoxyribose glycosyl bond of a damaged nucleotide residue, and leaves the sugar-phosphate backbone intact. The resulting abasic sites are then recognized by an apurinic/aprimidinic (AP) endonuclease that creates an incision on the backbone of the DNA at the 5' end of AP site (Sakumi and Sekiguchi, 1990). The completion of base excision requires removal of the 5 terminal deoxyribose-phosphate residue, which is catalyzed by the phosphodiesterase (AP lyase) activity of DNA polymerase β (Pol β) (Matsumoto and Kim, 1995). The resulting one-nucleotide gap is filled by DNA Pol β and sealed by either DNA ligase I or DNA ligase III with its accessory protein XRCC1 (Figure 3) in mammalian system (Lehman, 1998).

The BER pathway has been shown previously to depend strongly on the presence of nicotinamide adenine dinucleotide (NAD⁺). This dependency was proposed to be mediated through the catalytic activation of the nuclear enzyme poly(ADP-ribose) polymerase-1 (PARP-1) following its transient binding to BER-induced DNA strand break (Sato and Lindahl, 1992; Sato et al., 1993). PARP-1 is an abundant nuclear enzyme found in many eukaryotes, with the exception of yeast. This enzyme has high affinity for single- and double-strand DNA breaks. After binding to DNA strand breaks, PARP-1 catalyzes extensive synthesis of poly(ADP-ribose)




FIGURE 3. A model for base excision repair pathway. **AP endonuclease** = apurinic/apyrimidinic endonuclease; **POL.β** = polymerase β; **PCNA** = proliferating cell nuclear antigen; **FEN-1** = flap endonuclease-1

from NAD^+ and covalently modifies many nuclear proteins, including itself. The massive automodification of PARP-1 effects its dissociation from DNA strand breaks and inhibition of its catalytic activity. A modulatory role for poly(ADP-ribose) formation and PARP-1 automodification in BER has been proposed by Satoh and Lindahl (1992). According to this model, the unmodified enzyme binds tightly to DNA strand

interruptions formed either by ionizing radiation or through incision of AP sites during BER and interferes with the BER process because the bound PARP-1 molecule hinders access of the repair machinery to the lesion. Automodification and release of PARP-1 from the breaks allow the repair process to proceed. In the absence of NAD^+ or after the inhibition of poly(ADP-ribose) synthesis, PARP-1 persists on DNA breaks and

DNA repair is abrogated. Recently, it has been shown that the cells lacking PARP-1 have normal capacity to repair single-strand breaks inflicted by x-irradiation or breaks formed during the repair of modified bases (Vodenicharov et al., 2000).

Different DNA glycosylases (at least eight) are known to be present in human cell nuclei (Lindahl and Wood, 1999). These glycosylases remove different kinds of damage, and the specificity of repair pathway is determined by the type of glycosylase involved (Seeberg et al., 1995). The mode of action of these glycosylase and their three-dimensional structures have been described earlier (Cunnigham, 1997; Krokan et al., 1997; Parikh et al., 1998). These enzymes have a catalytic domain of ~250 amino acid residues and for additional interactions they also use their amino- and carboxy-terminal region (Lindahl and Wood, 1999). In general, these DNA glycosylases move along the minor groove of DNA until a specific type of damaged nucleotide is recognized. This enzyme then binds to the backbone of a damaged strand and slides out the damaged nucleoside residue to accommodate the damaged base in a specific recognition pocket and mediates cleavage (Parikh et al., 1998). Although these enzymes are essential for BER, in mice the knockouts of various DNA glycosylases have been shown to be viable (Wilson and Thompson, 1997), because the abasic sites are also known to be generated by nonenzymatic depurination.

Structurally, the AP endonuclease belongs to the superfamily of nucleases that also contains pancreatic DNaseI (Gorman et al., 1997). This enzyme flips out the base free deoxyribose residue from the DNA before chain cleavage and also recruits the Pol β to the site of repair (Bennett et al., 1997). The Pol β

contains two distinct domains, the larger domain is the polymerase domain, which helps in gap filling and the small basic amino-terminal domain that contain AP lyase (Phosphodiesterase) activity that excises the abasic sugar-phosphate residue at the strand break (Matsumoto and Kim, 1995; Sobol et al., 1996). A knock-out mutation of Pol β in mice has been shown to cause embryonic lethality (Gu et al., 1994). This finding suggests that either the single-patch mode of BER is essential for maintaining normal viability or that Pol β has an additional role in the cells such as in chromosomal DNA replication. The XRCC1 plays important role in bringing the Pol β and ligase III together, and it is also known to directly bind to the DNA single-strand break (Kubota et al., 1996; Marintchev et al., 1999).

Sometimes longer repair patches of 2 to 10 residues have also been observed (Matsumoto et al., 1994; Frosina et al., 1996). This might happen when the terminal sugar-phosphate residue has a more complex structure, which is resistant to cleavage by AP lyase, then the DNA strand displacement may occur. These longer repair tracts are thought to result from a nick translation reaction accompanied by strand displacement in the 5' to 3' direction, thereby generating a flap type of structure. The produced displacement flap structure is removed by flap endonuclease FEN-1 with the help of PCNA (Figure 3) (Harrington and Lieber 1994; Wu et al., 1996; Klungland and Lindahl, 1997). In this case the gap of few nucleotides long is filled by either Pol β or Pol δ (Fortini et al., 1998; Dianov et al., 1999) and sealed by DNA ligase I (Figure 3).

The BER in plants still has not been well studied when compared with mammalian systems, but this mechanism does

exist in plant (Talpaert-Borle and Liuzzi, 1982). During the early germination of *Zea mays*, the formation of AP site has been reported (Dandoy et al., 1987), which also suggested that plants do have BER pathway. A specific glycosylase enzyme (uracil-DNA glycosylase) was found in cultured cells of carrot that play a role in BER (Talpaert-Borle and Liuzzi, 1982). A cDNA clone encoding 3-methyl adenine glycosylase was reported from *Arabidopsis* and the encoded protein shown to contain glycosylase activity (Santerre and Britt, 1994). This gene was shown to be expressed in growing tissue in *Arabidopsis* (Shi et al., 1997). The gene encoding DNA ligase has also been cloned from *Arabidopsis* and sequence data are available in Genebank (Taylor et al., 1996b).

C. Nucleotide Excision Repair (NER)

NER is one of the most versatile DNA repair pathways operating in both prokaryotes and eukaryotes. Unlike other DNA repair pathways that are repair specific, NER is capable of removing various classes of DNA damage, including those induced by UV radiation (pyrimidine dimers) and chemicals (bulky DNA adducts) such as cisplatin, 4 NQO, benzo[a]pyrene, and aflatoxin (Lindahl and Wood, 1999; Balajee and Bohr, 2000).

The major difference between NER and BER is the way the DNA damage is removed. Basically, NER cuts out the damage as a part of an oligonucleotide fragment, while BER excises only one nucleotide. From 24 to 32 nucleotides can be removed by NER in eukaryote (Huang et al., 1992; Moggs et al., 1996). The repair of lesions over the entire

genome is called as global genome NER, while the repair of transcription blocked lesions present in transcribed DNA strands is called as transcription coupled-NER (TC-NER) (Mu and Sancar, 1997).


NER pathway in mammalian system involves product of at least 30 genes and most of them have been cloned (de Laat et al., 1999). In bacteria, mainly four proteins (UvrA, B, C, and D) carry out NER process. The UvrA, B, and C are required for incision reaction. The UvrA first dimerizes and then binds to UvrB, which recognize the damage on the DNA and causes local melting at 3' to the damage (Mazur and Grossman, 1991; Myles and Sancar, 1991). UvrB then makes a 3' incision (Lin et al., 1992), preceding the 5' incision by UvrC (Lin and Sancar, 1992). Incision is made at seventh nucleotide from 3' side and at fourth or fifth positions from 5' side of the damaged DNA (Sancar and Rupp, 1983). The UvrD helicase releases ~12 or ~13 bases long oligonucleotide containing the damage, DNA polymerase I fills the gap, and finally DNA ligase seals the repair patch. The NER processes in yeast and mammalian cells are very much alike. The molecular mechanisms of NER in a mammalian system have been described in detail in several reviews (Lehman, 1998; de Laat et al., 1999; Lindahl and Wood, 1999; Balajee and Bohr, 2000).

Most of the NER genes have been isolated through transfection of repair-deficient rodent mutant cell lines and the recovery of the excision repair cross complementing (ERCC) human genes. Nearly all ERCC proteins subsequently appeared to be involved in the human repair disorders *Xeroderma pigmentosum* (XP) and Cockayne syndrome (CS) and therefore are called XP and CS factors, respectively. The *XPA*, *XPB*, *XPC*, *XPD*, *XPF*, and *XPG* have been cloned from

the mammalian system. The various XP and ERCC factors take part in the different steps of NER. The NER genes isolated from human, yeast, and plant show significant sequence similarity, suggesting that the NER mechanism is conserved throughout the evolution (Hoeijmakers, 1993a,b; Xu et al., 1998). Unlike bacteria, yeast, and mammalian systems where NER has been studied extensively, in plants it is not well characterized yet. Although biochemical evidence suggests that such a DNA repair mechanism is present (McClennan, 1987; Britt, 1996;

Xu et al., 1998). Molecular characterization of gene products involved in DNA repair in plants has so far been limited to photolyases, the enzymes involved in photoreactivation (Ahmad et al., 1997), which is described later. Although NER has a very broad substrate specificity, it is primarily involved in the removal of UV-induced damage from the DNA. UV radiation induces the formation of two major UV photoproducts, the cyclobutane pyrimidine dimers and 6-4 photoproducts. The structure of these photoproducts is shown in Figure 4. These pyrimidine

A.



B.




FIGURE 4. Structure of two main UV-induced photoproducts. (A) A cyclobutane pyrimidine dimer (TpT). (B) 6-4 photoproduct (pyrimidine [6-4] pyrimidine dimer) (TpC). The dimers are formed between adjacent (5' and 3') bases on the same DNA strand.

dimers are highly toxic because they have effect on transcription. Their efficient removal is an essential function for any living cells that is exposed to sunlight. The cyclobutane pyrimidine dimers can be formed between any two adjacent pyrimidines (Figure 4A) (Protic-Sabljić et al., 1986; Doetsch, 1995). The 6-4 photoproducts are formed by covalent bond between the carbon 6 and carbon 4 of adjacent pyrimidines (Figure 4B). These are the most frequent occurring UV photoproducts and occur at 5'-T-C-3' (Figure 4B), 5'-C-C-3', and 5'-T-T-3' but not at 5'-C-T-3' sites in DNA (Doetsch, 1995). These photoproducts can be repaired by NER; enzymatic photoreactivation, recombination repairs, and by postreplications repair (Friedberg et al., 1995).

The molecular model for NER in mammalian system is shown in Figure 5. The NER sequentially involves recognition of DNA damage, dual incision on damaged strand, excision of damage containing oligonucleotide, DNA synthesis (gap-filling), and ligation to replace an excised oligonucleotide (Figure 5) (Wood, 1997; de Laat et al., 1999). The initiation of DNA repair is the first step in the NER. The XPC recognizes the damage and binds strongly to the damage and then recruits the entire repair protein apparatus to the damage (Drapkin et al., 1994; Masutani et al., 1994; Reardon et al., 1996; Sugawara et al., 1998). The XPC protein (125 kDa) exists in a complex with hHR23B (58 kDa), a human homolog of the *Saccharomyces cerevisiae* repair protein Rad23 that enhances XPC-dependent excision repair *in vitro* (Sugawara et al., 1998). The DNA damage is also recognized by the XPA protein in association with replication protein A (RPA) and forms a XPA-XPC-RPA complex on the dam-

age (Figure 5) (Matsuda et al., 1995; Saijo et al., 1996). The RPA is a heterotrimeric single-stranded DNA-binding protein that is required for both the incision reaction as well as repair synthesis (Coverley et al., 1992; Aboussekhra et al., 1995; He et al., 1995; Mu et al., 1995). RPA also binds to the nondamaged strand to prevent it from incision of the strand (which serves as a template for repair synthesis) and stabilizes the open structure (Kim et al., 1992; Seroussi and Lavi, 1993). RPA is also known to stimulate the interaction of XPA and ERCC1 (Li et al., 1994; Park and Sancar, 1994; Saijo et al., 1996), and on the other hand can bind to the endonuclease XPG (He et al., 1995) and possible XPF (Bessho et al., 1997). The XPA binds to basal transcription factor, TFIIF, and thereby recruits it to the damage area (Figure 5). The TFIIF is a 6-9 subunit complex containing XPB (ERCC3) and XPD (ERCC2) DNA helicases (Winkler et al., 1998; de Laat et al., 1999; Lindahl and Wood, 1999).

The action of TFIIF complex resulted in the formation of an open duplex structure around the damaged site that enables the structure-specific nucleases (XPG and ERCC1-XPF) for the dual incision (Evans et al., 1997; Mu et al., 1997a,b; de Laat et al., 1999; Lindahl and Wood, 1999). TFIIF and RPA can both bind to XPG, which makes the first incision while it cuts the damaged DNA strand on the 3' side of the damage (Figure 5) (O'Donovan et al., 1994; Matsunaga et al., 1995; Mu et al., 1996). The incision by XPG is made at 5 to 6 phosphodiester bonds away from the 3' side of the damage (Figure 5) (Svoboda et al., 1993). The XPG protein is a member of the FEN-1 family of structure-specific endonucleases and cuts at the junctions of du-




FIGURE 5. A model for nucleotide excision repair. The figure shows several steps, including damage recognition, unwinding of damage area, dual incision, excision of damage piece, gap filling, and the sealing (for details see the text).

plex and unpaired DNA (Lieber, 1997; de Laat et al., 1999). XPG is also known to interact with proliferating cell nuclear antigen (PCNA) (Gary et al., 1997), which is involved in DNA repair synthesis but dispensable for the incision stage of NER (Shivji et al., 1992). Next, the ERCC1-XPF complex makes the incision at 22 to 24 phosphodiester bonds away from the 5' side of the damage (Figure 5) (Svoboda et al., 1993; Mu et al., 1996). The dual incision is absolutely dependent on ATP hydrolysis (Svoboda et al., 1993). The specificities of these two nucleases make them ideally suited to cleave the opened up damaged site on either side of the damage. This results in the removal of a 24- to 32-residue oligonucleotides in length (Figure 5) (Huang et al., 1992; Moggs et al., 1996; Mu et al., 1997a,b; de Laat et al., 1999).

The final step in the NER pathway is the gap-filling or repair synthesis of the excised patch, which is performed by common DNA replication factors. An *in vitro* reconstituted repair reaction showed that efficient repair synthesis required the mammalian replication factors RPA, RFC, PCNA, and DNA polymerase δ or ϵ (Hubscher and Thommes, 1992; Shivji et al., 1995). RFC is needed to load PCNA onto the DNA (Podust et al., 1994), and PCNA is required for the initiation of repair synthesis. PCNA also stimulates polymerase activity. The newly synthesized repair patch, which exactly matches the excision patch, is then sealed by ATP-dependent DNA ligase, most likely ligase I (Figure 5) (Lindahl and Barnes, 1992; Petrini et al., 1995).

In plant, the NER system is still not well characterized, although biochemical evidence suggests that such a DNA repair mechanism is present (McCleannan,

1987; Britt, 1996; Vonarx et al., 1998; Britt, 1999). An UV-specific endonuclease resembling UVrABC nuclease, which is known to be involved in NER in bacteria, was reported and characterized from spinach (Doetsch et al., 1989). Molecular characterization of gene products involved in DNA repair in plants so far has been limited to photolyases (Ahmad et al., 1997). The UV products CPDs and (6-4) photoproducts are also known to be repaired in plant through a light-independent repair pathway that supports the existence of a pathway equivalent to NER mechanism in yeast and mammal (Howland, 1975; Quaitte et al., 1994; Taylor et al., 1996). NER was also suggested to occur in carrot protoplasts (Howland, 1975, Eastwood and McLennan, 1985), *Arabidopsis* and alfalfa seedlings (Pang and Hays, 1991; Quaitte et al., 1994), soybean chloroplasts and leaves (Cannon et al., 1995; Sutherland et al., 1996), and in different rice cultivars (Hidema et al., 1997). It was also reported in wheat leaf tissue (Taylor et al., 1996). The *Rad23* gene of yeast is involved in NER and transcription coupled repair (Mueller and Smerdon, 1996). Mutation of the yeast *Rad23* gene resulted in moderate UV sensitivity that was due to the defect in NER pathway (Verhage et al., 1996). In 1997 a structural homolog of *Rad23*, *OSRad23*, was reported from rice but its role in NER has not been determined (Shultz and Quatrano, 1997). Several putative *Rad23* homolog in *Arabidopsis* EST databases have been observed.

Until recently, the only evidence of the existence of light-independent DNA repair pathways in plants came from physiological studies based on dark repair of UV-induced pyrimidine dimers (Eastwood and McLennan, 1985; Quaitte et al., 1994; Taylor et al., 1996). In ad-

dition, UV-sensitive *Arabidopsis* mutants defective in dark repair have been described (Britt et al., 1993; Harlow et al., 1994; Jiang et al., 1997). Recently, a few plant homologs of some of the components involved in mammalian NER pathway have been reported. The molecular evidences for the conservation of the NER pathway came from the cloning of *Arabidopsis XPB* (*araXPB*) (Ribeiro et al., 1998) and the lily homolog of human *ERCC1* (Xu et al., 1998). The *araXPB* protein shared 50% identity and 70% conserved amino acids with yeast and human homologs. The plant *XPB* contained all the functional domains found in the other proteins, including nuclear localization signal, DNA-binding domain, and helicase motifs, suggesting that it might be playing a role in NER in plant cells (Ribeiro et al., 1998). However, the DNA unwinding activity of plant *XPB* has not been determined yet.

The plant *ERCC1* gene was isolated from male germline cells of lily, and the deduced amino acid sequence contained striking homology with human *ERCC1* and yeast *Rad10* (Xu et al., 1998). The lily *ERCC1* homologs were present in the genomes of taxonomically diverse plant species, including *Arabidopsis thaliana*, *Brassica napus*, *Zea mays*, *Oriza sativa*, *Nicotiana tabacum*, and *Lycopersicon esculentum*, as shown by Southern blot analysis of genomic DNAs (Xu et al., 1998). These findings suggested that the NER role of the *ERCC1* gene might be conserved in higher plants. The lily *ERCC1* expression was also reported to be upregulated in the male germline cells of plants (Xu et al., 1998). Pollen, as part of its developmental process, is exposed to solar UV radiation and other environmental mutagens after being released from the anther. This ex-

posure could inevitably lead to DNA damage in both vegetative and generative cell nuclei of pollen. The up-regulation of *ERCC1* homolog in lily generative cells suggested that a highly active DNA repair mechanism existed in male germline cells in order to protect germline DNA from heritable mutations resulting from damaged DNA (Xu et al., 1998). The protein encoded by the lily *ERCC1* gene was also able to correct the sensitivity to the cross-linking agent mitomycin C in *ERCC1*-deficient Chinese hamster ovary cells that further suggested that the NER mechanism is conserved in yeast, animals, and higher plants (Xu et al., 1998).

Recently, two different groups have reported the cloning of plant (*Arabidopsis*) homolog of human *XPF* and *S. cerevisiae* *RAD1* and they gave the name of the gene as *UVH1*, *Rad1* (Liu et al., 2000), and *AtRAD1* (Gallego et al., 2000). The *UVH1/Rad1* gene was cloned from *Arabidopsis* by positional cloning methods and has been mapped to chromosome 5 and also shown to be the product of a repair endonuclease (Liu et al., 2000). The *UVH1/Rad1* gene was shown to be strongly expressed in flower tissue as well as in other tissues (Liu et al., 2000). The *AtRAD1* gene from the other group is also shown to be involved in NER in the dark and is also shown to be ubiquitously expressed, which suggested that plant NER has some other roles (Gallego et al., 2000). Recently, the *XPF* gene was knocked out in tobacco by introducing *AtXPF* antisense in tobacco, which resulted in the increased UV sensitivity (Crockett et al., unpublished data). The transgenic tobacco plant carrying on *AtXPF* promoter *GUS* gene fusion showed relatively higher levels of *GUS* expression in generative cells of pollen and seeds. These findings suggested a key role of NER pathway in maintaining germline

DNA integrity and also in the removal of endogenous DNA lesions in seed, which accumulated during the seed aging process (Crockett et al., unpublished data).

D. Photoreactivation (PR) and Photolyases

It is clear now that UV radiation can damage plants, decreasing growth, and productivity (Teramura, 1983). The UV-induced DNA damages (CPDs and 6-4 pps) are removed or repaired by two different repair pathways. One is NER, which is described above, and other is PR which is a direct reversal phenomenon. PR is performed by the combined action of one or more proteins termed 'photolyases' and visible light in an error-free fashion (Sancar, 1994; Britt, 1999; Thoma, 1999). The photolyases specifically recognize and bind to the pyrimidine dimers form a complex that is stable in the absence of light. After absorbing a blue light photon, the pyrimidine dimers are reversed to pyrimidine monomers without excision of the damaged bases (Sancar, 1996; Thoma, 1999) as shown in Figure 6. Photorepair of CPDs have been reported in several plant species, including ginkgo (Trosko and Mansour, 1969), *Arabidopsis* (Pang and Hays, 1991; Britt et al., 1993), alfalfa (Quaite et al., 1994), soybean (Sutherland et al., 1996), cucumber (Takeuchi et al., 1996), rice (Hidema et al., 1997, Hidema et al., 2000), maize (Stapleton, 1992; Stapleton et al., 1997), and wheat (Taylor et al., 1996a).

Photolyases are known to be very specialized in terms of their substrate specificity (Britt, 1999). CPD photolyases were reported in bacteria, fungi, plants, inverte-

brates, and many vertebrates, while 6-4 photolyases were identified in silkworm, frog, fly, and rattlesnakes (Yasui et al., 1994, Sancar, 1996; Todo, 1999). The first higher eukaryotic photolyase was CPD specific and cloned from goldfish (Britt, 1999). Two distinct photolyases, one specific for CPDs and other specific for 6-4 products, were reported from *Arabidopsis* seedling (Chen et al., 1994). The two classes of CPD photolyases share only 10 to 15% sequence identity (Yasui et al., 1994). Class I is a microbial CPD photolyase and Class II is metazoan CPD photolyase. The *Arabidopsis* homolog of class II photolyase sequence (*PHRI*) corresponds to a gene (*UVR2*), which was identified via classic genetic analysis (Ahmad et al., 1997; Jiang et al., 1997). The phylogenetic analysis of photolyase homologs are described in Nakajima et al. (1998) and Britt (1999). Photolyases contain two prosthetic chromophores: FADH₂ and either methenyl tetrahydrofolate or 8-hydroxy-5-deazaflavin (Sancar, 1994). The second chromophore works as a light-harvesting antenna. The wavelength of light for photoreactivation ranges from the visible (500 nm) to the UV-B (300 nm). The existence of PR in plant was first reported by Ikenaga and Mabuchi (1966), who demonstrated that the frequency of endosperm mutations generated by UV radiation of maize pollen dropped substantially if the pollen was exposed to visible light after UV irradiation. A photolyase cDNA was reported earlier from white mustard (Batschauer, 1993). The crystal structures of CPD photolyase of *E. coli* and *Anacystis nidulans* have been resolved. This suggested that the photolyases flip the pyrimidine dimer out of the duplex into the hole that contains the catalytic cofactor (Park et al., 1995; Tamada et al., 1997) followed by splitting the cyclobutane ring after light absorption.




FIGURE 6. Schematic illustration of photoreactivation. Photolyase binds to pyrimidine dimer (T = T), split the dimer and restore it to native pyrimidine monomer in a light-dependent reaction.

The plant blue light photoreceptor genes were discovered during the photolyase research and named *CRY* after cryptochrome. The *CRY* of plant and animal have some sequence homology with DNA photolyase (Todo, 1999), but they do not contain any repair activity. In yeast it is known that repair by photolyase is faster than NER especially in nucleosome-free regions, for example, promoters and origin of replication re-

gions (Thoma, 1999). The photolyase and NER can also play complementary roles in the genes that are transcribed by RNA polymerase II; they remove the damage in the nontranscribed and transcribed strands, respectively (Livingstone-Zatchej et al., 1997; Suter et al., 1997; Thoma, 1999). Aboussekhra and Thoma (1999) have shown that NER can remove CPDs, which are not accessible to photolayase, such as CPDs gen-

erated in the TATA-box of the yeast *SNR6* gene.

Increased sensitivity to UV radiation may result from failure to repair photodamage in DNA. The UV radiation-sensitive *UVR1* mutant of *Arabidopsis* cannot photorepair 6-4 photoproducts (Britt et al., 1993). Landry et al. (1997) showed that another *Arabidopsis* UV radiation-sensitive mutant, *UVR2-1*, was deficient in photorepair of CPDs and cloning and sequencing of the photolyase genes from UV radiation-sensitive and wild-type strains indicated alterations of DNA sequence that led to defective photolyase protein (Ahmad et al., 1997). These data suggested that such deficiencies might be ameliorated by restoration of photolyase function in photorepair-deficient strains. However, for such restoration exact knowledge of the molecular origin of their photorepair deficiency is essential, as suggested by Hidema et al. (2000). It is important to know whether the deficiency is in the photolyase protein itself, or it has resulted from other cause, such as a regulatory mutation that causes the production of fewer normal photolyase molecules per cell. The UV-B radiation-sensitive rice cultivar Norin 1, a progenitor of many Japanese commercial rice strain, has been reported to be deficient in photorepair of CPDS (Hidema et al., 1997). Recently, the same group, Hidema et al. (2000), reported that the photorepair deficiency in Norin 1 is due to the alteration in the structure of the photolyase enzyme rather than a regulatory mutation. Their findings indicated that a strategy of increasing the activity of photolyase through selective breeding or engineering could increase resistance to UV radiation and perhaps the productivity of such cultivars (Hidema et al., 2000).

E. Bypass Damage Repair and DNA Polymerases

It is known that the pyrimidine dimers can act as blocks to DNA replication, so cells have to possess specialized low fidelity and often error-prone DNA polymerases that can bypass such lesions and promote replication of damaged DNA (Johnson et al., 1999c). Usually at cell-cycle checkpoints, these damages are repaired before replication or cell division proceeds. If the damage is not repaired properly, then the cell cycle checkpoints can lead to the inhibition of cell cycle progression (Costanzo et al., 2000). In yeast *S. cerevisiae* the catalytic subunit (Rev3) of DNA polymerase ζ together with an accessory factor Rev7 can bypass pyrimidine dimers and other adducts in DNA (Lawrence and Hinkle, 1996). Some other polymerases, for example, yeast, Rev1 (hREV1 in human), *E. coli* UmuD'2C (*E. coli* Pol V), *E. coli* DinB (Pol IV), are also known for bypassing the damage before the action of replicative polymerases (Nelson et al., 1996; Gerlach et al., 1999; Lin et al., 1999; McDonald et al., 1999; Tang et al., 1999; Wagner et al., 1999). The DNA polymerase encoded by yeast *RAD30* and human *hRAD30A* also bypasses a cis-syn thymine-thymine dimer efficiently and accurately (Johnson et al., 1999 a,b; Masutani et al., 1999; Washington et al., 1999; Johnson et al., 2000b, Washington et al., 2000). The damage bypassed by polymerases that insert wrong residues result in mutations, which can create more problems.

The human gene *hRAD30B* (McDonald et al., 1999), which is a analog of yeast *RAD30B*, encodes the ninth eukaryotic DNA polymerase named Pol ι , which misincorporates deoxynucleotides at a high

rate (Johnson et al., 2000 a). In general, DNA polymerases extend from mispaired bases at about the same frequency at which they insert the respective mispair (Goodman et al., 1993). A new model for mutagenic bypass of DNA lesions in eukaryotes showed that two DNA polymerases (τ and ζ) act sequentially (Johnson et al., 2000 a). Pol τ functions in damage bypass by inserting deoxynucleotides opposite DNA lesions, whereas Pol ζ acts at the subsequent step of extending from them (Gibbs et al., 1998; Johnson et al., 2000a).


Whether the bypass repair is responsible for mutagenesis in plants remains to be studied in detail. However, in maize pollen the UV-induced mutations were found to be nontransmissible, which suggested that the mutations are of deletion type rather than point mutation (Nuffer, 1957).

F. Double-Strand Break (DSB) Repair

As early as in 1931, by using cytological tools, Barbara McClintock showed that the outcome of radiation-induced chromosome breaks is translocations, deficiencies, ring chromosomes, and end fusions (McClintock, 1931). She suggested that the broken ends can fuse with the same or other broken ends or chromosomes, we now call this DSB repair. DSBs are usually created by ionizing radiation (Dizdaroglu, 1992), chemical agents, during meiosis (Friedberg et al., 1995) or transposition reaction (Lieber, 1991) and also as a result of a nick in the single-stranded region due to mechanical stress of a chromosome (Britt, 1999). The unprotected DSB is a dangerously cytotoxic lesion. If not repaired immediately, DSBs can

be attached to by unwelcome recombination events, which can lead to loss of portions of chromosomes or to rearrangements (error prone). Usually DSBs are repaired by either homologous recombination or by nonhomologous end-joining (NHEJ), which involves the joining of two broken ends (Britt, 1999). Homologous recombination is particularly effective in S phase when the break can be repaired using genetic information from a sister chromatid, whereas NHEJ is thought to be effective at all times in the cell cycle (Takata et al., 1998; Essers et al., 2000). NHEJ also plays an important role in DSB repair during V(D)J recombination in antibody coding fragments (Blunt et al., 1995). The model of DSB is shown in Figure 7. In general, the organisms with relatively compact genomes (yeast, bacteria, etc.) repair the DSB via homologous recombination, provided some homology is available and organisms with larger genomes follow NHEJ (Britt, 1999). In plants the DSB repair products have been characterized as excision products of transposable elements, or insertion products of *Agrobacterium* T-DNA (Ohba et al., 1995; Scott et al., 1996).

Recombination is initiated by a 5' to 3' exonuclease (Figure 7) that generates long single-stranded 3' tails (White and Haber, 1990; Sugawara and Haber, 1992). In homologous recombination, one of these tails invades a homologous DNA duplex. Overall, this resulted in a formation of Holliday Junction, which resolved and forms two recombinant DNA molecules with or without crossing over. The DSB repair by homologous recombination with another allele can be achieved with high fidelity, whereas repair by NHEJ may result in lost or changed genetic information.



Nonhomologous recombination Homologous recombination

FIGURE 7. Models for double-strand break repair. In these schematic models, double-strand breaks caused by either irradiation or chemical assault or during meiosis and transposition reactions or due to mechanical stress can be repaired by nonhomologous end joining or nonhomologous recombination or homologous recombination.

In plants, the DSB repair is still poorly understood. Evidence from transformed plants suggests that the majority of transformed DNAs are integrated via NHEJ. However, plants are capable of repairing DSB via either pathway. In plants a few genes that are involved in DSB repair have been isolated. The homologs of RecA genes (*DMC/LIM15/RAD51*) have been reported from plants (Kobayashi et al., 1993; Klimyuk and Jones, 1996; Doutriaux et al., 1998). The RecA-like recombination protein was shown to be localized on chromosomes of the lily at various meiotic stages (Terasawa et al., 1995). The DSBs in plant somatic tissues can be repaired by homologous recombination. Greater induction of homologous recombination was observed when site-specific DSBs were created using transposon or rare-cutting endonucleases. The first molecular data on end-joining in plants came from the analysis of footprints left behind after transposable element excision (Rinehart et al., 1997). These footprints usually consist of minor changes (i.e., a few base pair deletions, or inversion), and the exact configuration of DNA ends produced by transposon excision remains unknown. Recently, a Rad50 homologue gene and the encoded protein have been isolated from *Arabidopsis* and shown to be involved in meiosis and DSB repair in plants (Gallego et al., 2001).

The repair of DSBs by NHEJ requires the products of the *XRCC4*, *XRCC5*, *XRCC6*, and *XRCC7* genes in mammalian system (reviewed by Chu, 1997; Critchlow and Jackson, 1998). *XRCC4* encodes a protein (XRCC4) that forms a heterodimer with ligase IV that rejoins two DNA ends in the last step of V(D)J recombination and NHEJ to repair DSBs. *XRCC4*-defective cells are extremely sensitive to ionizing radia-

tion and disruption of the *XRCC4* gene or ligase IV results in embryonic lethality in mice (Barnes et al., 1998). Recently, the crystal structure of *XRCC4* has been resolved at 2.7 Å, which suggests a dumbbell-like tetramer structure (Junop et al., 2000). *In vitro*, *XRCC4* stimulates adenylation of ligase IV, the first chemical step in DNA ligation (Modesti et al., 1999). *XRCC4* may also bridge two ends for ligation, perhaps by interacting with Ku protein already present at the ends (Chen et al., 2000). *XRCC5* and *XRCC6* encode the 70- and 80-kDa subunit of Ku, and *XRCC7* encodes the catalytic subunit of DNA-dependent protein kinase (DNA-PKcs). In mammalian system, it has been shown that DNA-PK is involved in the major pathway for repairing DSB (Jeggo et al., 1995). DNA-PK is a heterotrimer consisting of a 450-kDa catalytic subunit (DNA-PKcs) and Ku80 and Ku70 regulatory subunits (Anderson, 1993; Tuteja et al., 1994, Tuteja and Tuteja, 2000). The Ku80 and Ku70 of DNA-PK bind to DNA ends (broken ends), which activate the kinase activity of DNA-PKcs. The DNA-PKcs is a serine/threonine protein kinase and can phosphorylate a variety of substrates *in vitro* (Tuteja and Tuteja, 2000). Finally, the Ku80 and Ku70 bring the DNA broken ends together and then DNA ligase joined the ends (Figure 7). The end-binding activity of Ku serves as an entry site for other proteins and the phosphorylation activity of DNA-PKcs activate other repair factors (Tuteja and Tuteja, 2000). DNA-PK is the only factor known to participate in NHEJ. DNA-PKcs is a member of the phosphatidylinositol 3-kinase (PI3K)-related kinase family, as are ATM and ATR. All three proteins exhibit a strong sequence homol-

ogy to the PI3Ks, especially in the catalytic core domain that binds and phosphorylates the phosphoinositol head group of phosphatidylinositol. However, no phosphorylation of lipid substrates has been observed by DNA-PKcs or by other members of this PI3K-related family of kinases (Carpenter and Cantley, 1996; Wymann and Pirola, 1998).

The mutants deficient in Ku or DNA-PKcs are deficient in the repair of DSBs and also sensitive to lethal effects of ionizing radiation. A new activity of Ku as an unwinding enzyme was also reported (Tuteja et al., 1994; Tuteja et al., 1997). The analog of Ku in plant has not been reported yet. However, we have shown through Western blotting that the Ku or Ku-like protein are also present in plant (Tuteja and Tuteja, 2000). However, by using human Ku antibodies and pea cDNA library, we got the positive clone that later turned out to be nonspecific and actually was the clone of calnexin, a Ca^{2+} -binding molecular chaperone protein (Ehtesham et al., 1999).

In plants NHEJ has been analyzed in a more direct and thorough manner. Linearized plasmid DNA has been introduced into tobacco cells, and novel joints that formed between the plasmid ends have been sequenced (Gorbunova and Levy, 1997). In another study, DSBs were induced at an I-SceI recognition site, within a negative selectable marker genome on a chromosome, and new junctions were sequenced (Salomon and Puchta, 1998). Gorbunova and Levy (1997) summarized the results of these studies as follows:

- End joining by simple ligation with no sequence alteration is rare.

- End-joining is usually associated with deletions ranging from 1 bp up to > 1 kb.
- Rejoining frequently occurs at short repeats.
- A distinct feature of the junctions is the presence of relatively large insertions, up to 1.2 kb long.

The filler DNA could consist of a simple insertion corresponding to a plasmid or genomic sequence. Insertions might also have a complex structure consisting of a patchwork of different plasmid and tobacco genomic sequences.

In response to DSB, different repair routes can be taken. The break can also be simply ligated, without any modification of the ends or different events can happen at the broken site, such as exonuclear degradation or exposure of 3'-ss end, which was proposed to be the first step of DSB repair pathway in plants. When annealing of the exposed ends occurs at sites of microhomology, as often found in plants (Gorbunova and Levy, 1997; Salomon and Puchta, 1998), this results in a deletion of the region between the microrepeats. This process is reminiscent of the single strand annealing (SSA) mechanism, which was also proposed for extrachromosomal homologous recombination events in plants (Puchta and Hohn, 1991). The outcome of DSB repair is also the presence of templated filler DNA at the repair site (Gorbunova and Levy, 1997). Formation of such filler DNA can be best explained by the synthesis-dependent strand annealing (SDSA) mechanism that involves template invasion and DNA synthesis (Gorbunova and Levy, 1999). The SSA and SDSA appear to be the two prominent mechanisms for DSBs repair in plants. They can be used either for homologous or nonhomologous recombination.

G. Mismatch Repair (MMR)

Because of the errors of replication or homologous recombination, mismatched bases can be produced (Friedberg et al., 1995; Friedberg and Wood, 1996). The MMR basically discriminates between the correct and incorrect bases and finally corrects the error after DNA synthesis. Most of the present understanding of eukaryotic MMR has come from studies of the eukaryotic homologs of the bacterial MutS and MutL proteins (Kolodner, 1996; Kolodner and Marsischky, 1999). The model of MMR in *E. coli* is shown in Figure 8. In *E. coli* the MutS dimer recognize the mispairs and then bind on it (Grilley et al., 1989) followed by MutL binding, which activates MutH (endonuclease) that makes a single-strand incision. This incision by MutH can be on either side of the mismatch (Au et al., 1992). Subsequently, the excision initiates at the incision (nick) and proceeds toward the mismatch (Grilly et al., 1993). To fill the gap (100 to 1000 nucleotide gap) the original template strand can then be replicated and finally sealed by ligation (Figure 8). The proteins involved in the last step of eukaryotic MMR are DNA pol δ , RP-A, PCNA, and RFC (for a review see Kolodner, 1996; Kolodner and Marsischky, 1999).

The homologs of MutS and MutL have been reported in yeast and human (Modrich, 1994; Drummond et al., 1995; Palombo et al., 1995; Hunter et al., 1996; Johnson et al., 1996; Palombo et al., 1996). In plant, the homolog of MutS was reported from *Arabidopsis* (Culligan and Hays, 1997), but the function has not been determined yet.

IV. ROLE OF DNA HELICASES IN DAMAGE AND REPAIR

The DNA helicases are ATP-dependent DNA unwinding enzymes and thus play an important role in repair as well as in replication, recombination, and transcription (Tuteja and Tuteja, 1996; Tuteja, 1997). The unrepaired DNA lesions can affect the outcome of all these processes. The helicases are among the first proteins that would encounter DNA damage during all the processes above. The interstrand or intrastrand DNA lesions can block or impair the DNA helicase action. The XPB and XPD are repair helicases that are required for local unwinding of the damage area of duplex DNA in NER in the mammalian system (Schaeffer et al., 1993; Schaeffer et al., 1994). During DNA repair, especially in NER, the damaged area on DNA has to unwind first in order to get proper repair, because most of the DNA repair machinery enters through single-stranded DNA. This single-stranded DNA is provided transiently by the DNA repair helicase(s). This unwinding process by DNA helicases is an energy-requiring process that comes through the hydrolysis of ATP by the intrinsic DNA-dependent ATPase activity of the helicase protein itself (Tuteja, 2000).

DNA helicases generally bind to ssDNA or ss-/ds-DNA junctions and translocate unidirectionally along the bound DNA into the duplex region, disrupting the hydrogen bonds linking the two strands (Tuteja and Tuteja, 1996). The XPB can unwind DNA in the 3' to 5' direction (Schaeffer et al., 1993; Roy et al., 1994) similar to previously reported human DNA helicases (HDH) I, II, III, V, and VI (Tuteja et al., 1990; Tuteja et al., 1992 to 1995), pea chloro-




FIGURE 8. A model for mismatch repair in *E. coli*. The eukaryotic process is believed to share many of the features shown here.

plast DNA helicases I and II (Tuteja et al., 1996; Tuteja and Phan, 1998), pea nuclear DNA helicase 45 (Pham et al., 2000), and pea nuclear DNA helicase 65 (Tuteja et al., 2001). However, XPD unwinds the DNA opposite to XPB polarity, that is, in 5' to 3' direction (Sung et al., 1993), which is similar to HDH IV (Tuteja et al., 1991). In order to better understand NER in plant, it is important to isolate the DNA repair helicases

along with other components. These repair helicases could be analogs of XPB and XPD that need to be cloned in an expression vector for purification of the encoded proteins. Furthermore, these proteins need to be characterized in detail. XPC-hHR23B may assist this strand separation by interacting with TFIIH (Drapkin et al., 1994). The XPB and XPD both contain the seven conserved domains (helicase motifs) of the super-

family of DNA and RNA helicases (Gorbalenya and Koonin, 1993; Tuteja, 2000). However, the sole presence of helicase motifs in the protein may not prove that it is a helicase. For example, an *E. coli* Mfd protein (a strand-specific repair protein) (Selby and Sancar, 1993) and a human Rad54 protein (Swagemakers et al., 1998) both contain conserved helicase motifs but lack the unwinding activity. So a direct biochemical demonstration of ATP-dependent DNA or RNA unwinding activity is essential before a protein is classified as helicase (Tuteja, 2000).

In *E. coli* the Uvr₂B and UvrD are the repair DNA helicases involved in NER. The DNA unwinding activity of UvrD helicase was shown to be exquisitely sensitive to major groove DNA intercalators (George et al., 1992), while it was moderately sensitive to UV radiation. In general, the UV-induced lesions or other helix-distorting intrastrand adducts are better tolerated by DNA helicases than those produced by intercalating agents. However, in yeast *S. cerevisiae* the Rad3 protein, a repair helicase, is absolutely required in NER. It is found to be inhibited by UV damage and intrastrand cisplatin adducts located on the strand along which the enzyme translocates, whereas lesions on the opposite strand had no effect (Villani and Tanguy Le Gac, 2000). Calf thymus DNA helicase E is a 3' to 5' helicase that can unwind DNA duplex from nicks and has been suggested to play a role in DNA repair (Turchi et al., 1992).

Ku protein is another DNA helicase (human DNA helicase II) that plays an important role in double-strand break repairs (Tuteja et al., 1994; Tuteja and Tuteja 2000). Ku is a heterodimeric protein composed of ~70 and ~80 kDa subunits (Ku 70 and Ku 80) originally iden-

tified as an autoantigen recognized by the sera of patients with autoimmune diseases. It is an abundant nuclear protein found in vertebrates, insects, yeast worms, and also in plant (Tuteja and Tuteja, 2000). Ku contains ssDNA-dependent ATPase, DNA end binding, and ATP-dependent DNA unwinding activities (Tuteja et al., 1994). It prefers to unwind the partially, unwound DNA substrates and moves in the 3' to 5' direction on the bound strand (Tuteja et al., 1994). Ku also functions as a regulatory subunit of DNA-PK that phosphorylates many proteins, such as p53, SV-40 large T antigen, RP-A, hsp90, c-jun, C-Fos, Sp-1, C-Myc, TFIID, etc. (Tuteja and Tuteja, 2000). The DNA unwinding activity of Ku was found to be severely inhibited by DNA intercalating agents such as nogalamycin, daunorubicin, actinomycin, and ethidium bromide (Tuteja et al., 1997).

In general, intrastrand DNA damage seems to affect DNA unwinding activity of the helicase *in vitro* when placed on the strand to which the protein translocates with exception to the SV-40 T antigen and RecA helicases (Villani and Tanguy Le Gac, 2000). The DNA helicases involved in initial stages of NER are more sensitive to DNA damage (e.g., UvrAB, Rad3, or XPD) when compared with the helicases that are involved in the subsequent removal of the damaged DNA (e.g., UvrD).

V. DESSICATION-RELATED DAMAGE, TOLERANCE, AND DNA REPAIR

Plant seed, the product of fertilization, during its final stages of development and maturation undergoes desic-

cation, resulting in a significant reduction in its water content. In addition, metabolic activities are also reduced to undetectable levels, leading seed embryo in quiescent state. Under low temperature and moisture, seed can now be stored for long periods until it is reactivated by favorable seed germination conditions. Seed viability loss has been shown to result in a loss of membrane integrity, impaired RNA, and protein synthesis and ATP production (Priestley et al., 1985; Bailly et al., 1996). Ultrastructural, biochemical, and biophysical studies have shown that the loss of desiccation tolerance is correlated with an irreversible alteration to the physical properties of membrane lipid bilayers, increased liquid crystalline-gel phase transition temperature, deesterification of phospholipids, and an accumulation of several products of peroxidized lipids. Observations that lipid-rich seeds tend to have a limited longevity tend to suggest that seed aging based on lipid peroxidation is important (Priestley et al., 1985).

Free radicals and lipid peroxidation are considered to be major cause of seed deterioration (Leprince et al., 1996). Ungerminated seeds and pollen are known to contain and accumulate free radicals during artificial aging (Priestley et al., 1985) and germinating seeds after dehydration (Senaratna and McKersie, 1986). Changes in lipid peroxidation and in the molecular defenses against the activated forms of oxygen have also been observed in dry stored seeds and seed after accelerated aging. The loss of seed viability has been correlated with the MDA (malondialdehyde) content, a product of lipid oxidation. Reuzeau and Cavalie (1995) found a significant difference in the MDA content of seeds able or unable to germinate. A higher

lipid peroxidation was found in seeds unable to germinate. It has been suggested that secondary toxic products produced as a result of oxygen-dependent reactions in imbibed seeds by the accumulation of oxygenated fatty acids could participate in altered cellular components leading to a decrease in seed viability (Gidrol et al., 1989; Reuzeau and Cavalie, 1995). When such seeds are hydrated, toxic oxygen products, if they are not efficiently detoxified, can allow damaging reactions and prevent seed germination. Because dry seeds are affected in their ability to detoxify O₂-related toxic products, it seems reasonable to suggest that important changes in soluble and membrane entities might interfere in the signal transduction necessary for the initiation of germination process (Reuzeau and Cavalie, 1995). Similar conclusions on the role of lipid peroxidation were drawn when accelerated aging of seed was studied (Bailly et al., 1996). Nonviable seeds with accelerated aging induced the accumulation of malondialdehyde, indicating that seed deterioration was associated with lipid peroxidation. No direct correlation between lipid peroxidation and deterioration in membrane integrity was found. However, loss of seed viability was associated with a decrease in superoxidase dismutase, catalase and glutathione reductase (GR) activities. This study established a clear relationship between the loss of detoxifying enzymes such as catalase, and GR, lipid peroxidation, and seed deterioration during accelerated aging. Whether oxidative damage and decrease in SOD, catalase, and GR activities are the cause or consequence of the loss of seed viability was not established (Bailly et al., 1996).

The desiccation tolerance mechanisms allow cells to withstand the loss

of the major percentage of its water content and to survive in a dry state for prolonged periods. The seeds, spores, wind-dispersed pollen, and specialized 'resurrection' species can withstand desiccation without incurring cellular damage, and they naturally dehydrate to moisture contents that will result in cell death in other plant tissues.

The role of free radicals in loss of desiccation tolerance in germinating seed has also been studied (Leprince et al., 1990). Desiccation of 24 h during germination resulted in a significant increase in lipid peroxidation in the dehydrated tissues, and rehydration not only fails to reduce lipid peroxidation but resulted in a 15-fold increase in MDA accumulation, indicating that desiccation treatment results in irreversible peroxidative damage to lipids. It is not clear whether oxygen radicals are the cause of or a result of the damage following desiccation, but it is well established that desiccation alters the properties of membranes (Senaratna and McKersie, 1986). A desiccation-induced rise in lipid peroxidation originates from oxygen radicals and a loss of desiccation tolerance is a consequence of increased formation of one or more forms of activated oxygen coupled with decreased protection from SOD and peroxidases. It is now clear that damages to DNA are harmful, which is why the repair of DNA damage is of vital importance to the cell in order to keep its normal function.

A. In Bacterial Spores

Desiccation tolerance of bacterial spores depends on the conversion of B-form DNA to a more condensed A-

form (Setlow, 1992a,b). The DNA in dormant spores of *Bacillus* species is associated with alpha/beta-type, small acid-soluble proteins (SASP), which are double-stranded DNA-binding proteins whose amino acid sequence has been highly conserved in evolution. The expression of these proteins occurs at a specific stage of spore maturation when the B-form DNA duplexes lose water and are converted to A-form conformation (Setlow, 1992a,b). These proteins bind to A-form, conferring stability on the DNA during the final stages of dehydration and through the dehydrated stage. These proteins are released by the action of the sequence-specific protease (termed GPR) that degrades SASP during germination. These spore protease of *Bacillus* are synthesized during sporulation as an inactive precursor termed P46. The activation of this protease during rehydration leads to the removal of SASPs and the return of the DNA to the B-form (Illadas-Aguar and Setlow, 1994). The saturation of spore DNA with SASP provides protection against DNA base loss due to wet heat (i.e., de-purination) and breakage of DNA single-strands by desiccation and hydrogen peroxide. The spores lacking a/b SASP are much more sensitive to desiccation, dry and wet heat, and hydrogen peroxide than are wild-type spores. All these treatments have been shown to kill spores in large part by damaging DNA (Setlow and Setlow, 1996). Binding of a/b type SASP also helps in conferring resistance to UV damage (Setlow and Setlow, 1996).

The major deleterious DNA photoproduct generated by UV irradiation of growing cells is a cyclobutane-type dimer between adjacent thymine residues (termed TT), but TT is not formed

in spores. The major UV photoproduct formed in spore DNA is a thymine-thymine adduct termed the spore photoproduct (SP), which is also formed between adjacent thymine residues. SASP deficient spores are ~30 times more UV sensitive than the wild-type spores, and UV generates a large amount of TT in deficient spores. According to Setlow and Setlow (1996), the generation of TT is the cause for the increased UV sensitivity of mutant spores. These authors examined the expression of genes involved in DNA repair during germination and outgrowth of both wild type and a-b-spore, which were exposed to a variety of treatments. This study has shown that the repair of DNA damage during spore outgrowth is an important component of spore resistance to a number of DNA-damaging treatments. The SASP-type proteins have not been identified in plants as such. However, other proteins may also play an important role in survival of dehydrated seeds and pollen. Dehydrins and LEA-type proteins may have some role in desiccation tolerance.

The *Deinococcus radiodurans* is a non-spore-forming bacterium that exhibit a remarkable capacity to resist the lethal effects of ionizing radiation. It has been reported that *D. radiodurans* ionizing radiation resistance is incidental, a consequence of this bacterium's adaptation to dehydration (Mattimore and Battista, 1996). The authors showed that *D. radiodurans*'s cellular responses to ionizing radiation and dehydration significantly overlap by showing that 41 ionizing radiation-sensitive strains of *D. radiodurans* were also sensitive to desiccation. It was also inferred that loss of DNA repair ability in the ionizing radiation-sensitive strains was responsible for the desiccation sensitivity of these strains.

B. In Seeds

It has been known for more than 2 decades that a feature of loss of viability in stored dry seed is the increasing loss of integrity of nuclear DNA with increasing levels of low-molecular-weight oligonucleotide fragments (Cheah and Osborne, 1978; Elder et al., 1987; Elder and Osborne 1993; Leprince et al., 1996). When rye embryos that are nonviable are imbibed for 2.5 h the extent of fragmentation of DNA shows a significant increase pointing toward the continuous activity of endonuclease cleavage in the absence of DNA repair (Cheah and Osborne, 1978; Elder et al., 1987). Embryos of imbibed dormant *A. fatua* also showed an efficient repair of radiation-induced lesions in the DNA of dry seeds, and this DNA repair also remained stable when embryos were dried back. On the basis of results from various experiments, Boubriak et al. (1997) have suggested that the failure to restore genetic integrity to the genome on rehydration of an embryo whose DNA is desiccation damaged could be a major factor in determining desiccation tolerance. The ability to repair DNA when embryos are rehydrated is critical to maintaining a transcriptionally functional genome.

Elder et al. (1987) have shown accumulation of single- and double-strand breaks in the embryos of seeds during the loss of viability in the dry state. In contrast to hydrated cell, which can effectively repair DNA damage, the nuclei of dry cells have to accumulate DNA damage until rehydration activates DNA repair machinery. The stability of the DNA on desiccation and the capacity for repair of DNA on rehydration have been suggested as integral components

of a total desiccation tolerance mechanism.

C. DNA Integrity in Pollens

Osborne and Boubriak (1994) have shown that dry birch pollen (14% moisture content) can be fully hydrated in moist air and dehydrated back to the original dry weight without the loss of germinability and without the fragmentation damage detectable in the extracted DNA. At the time of anther dehiscence, birch pollen is haploid and bi-nucleate and arrested at the G2 stage of the cell cycle. The authors addressed the question whether there is evidence for DNA repair during the air hydration of birch pollen. Pollen that has been exposed to increasing durations of hydration in moist air prior to the transfer to the liquid germination medium showed similar levels of H3-methyl thymidine. It was suggested that an excision repair measured as unscheduled DNA synthesis does not occur during the relatively low level of hydration possible in moist air. As pollen gets exposed naturally to UV in solar radiation following release from the anther, it is the likely candidate for UV-induced heritable mutations.

The role of DNA repair in the desiccation tolerance of pollen and spores is not unlike that ascribed to DNA repairs in bacterial spores (Setlow, 1992a,b). In their dehydrated state, pollen and seed continue to accumulate DNA damage that cannot be repaired until the rehydration occurs. The low water content in the desiccated stage prevents enzymatic activity and therefore DNA repair. No evidence of DNA repair has been observed in dry pollen or seed. Thus DNA damage cannot be repaired until pollen

and seed are rehydrated. Given the DNA damage that may accumulate in pollen and spores during the dormant period, they may contain high levels of enzymes needed for DNA repair.

Jackson and Linskens (1978) made comprehensive analysis of DNA repair system in *Petunia* pollen. The mature pollen of *Petunia* carries generative nucleus blocked at G2 stage of cell cycle. The addition of 4-nitroquinolide 1-oxide to the liquid germination medium led to unscheduled DNA synthesis in pollen. Mature *Petunia* pollen is bicellular and thus has two nuclei, that is, generative and vegetative nuclei. In these studies on *Petunia* pollen it was not feasible to discern whether the repair synthesis takes place in either or both nuclei of pollen. It was shown that unscheduled DNA synthesis occurs after ultraviolet radiation. As this unscheduled DNA synthesis was not affected by hydroxyurea, an agent used to discriminate between replicative and repair synthesis, the authors suggested that dark repair of DNA takes place in germinating pollen of *Petunia* (Jackson, 1987).

VI. DNA REPAIR GENES IDENTIFIED BY SEQUENCE HOMOLOGY

Recently, Wood et al. (2001) have described a catalog of 130 known human repair genes on the basis of function or sequence homology to known repair genes. Here we are mainly describing the plant repair genes found on the basis of sequence homology. The completion of the first plant genome sequence of *Arabidopsis thaliana* is a landmark of twentieth century in plant research (The *Arabidopsis* Genome Ini-

tative, 2000). The information it contains will certainly give a boost to every area of plant science, including DNA repair in near future. *A. thaliana* is a modest little flowering brassica plant chosen first for plant genome sequence because it has many advantages for genome analysis, including a small size, large number of offspring, a short generation time, and a relatively small nuclear genome. In 1977 William Curtis, a British botanist and apothecary, described *A. thaliana* in his flora landinensis as a plant of “no particular virtues or uses”, but now with the report of its genome sequence it becomes the most significant plant in the kingdom (Dennis and SurrIDGE, 2000). The sizes of the *Arabidopsis* genome of nuclear, plastid and mitochondrial are 125 Mb, 560 kb, and 367 kb, respectively. It has five chromosomes (chromosomes 1 to 5) with respective sizes of 29.1, 19.6, 23.2, 17.5, and 26.0 Mb. The evolution of *Arabidopsis* involved a whole-genome duplication, followed by subsequent gene loss and extensive local gene duplications. The nuclear genome contains 25,498 gene encoding proteins from 11,000 families, which is the largest gene set published to date (The *Arabidopsis* Genome Initiative, 2000).

Comparing the *Arabidopsis* genome with other species (Eisen and Hanawalt, 1999) indicated that *Arabidopsis* has a set of DNA repair and recombination (RAR) genes similar to most other eukaryotes. The pathways represented include PR, DNA ligation, NHEJ, BER, MR, NER, and many aspects of DNA recombination. The *Arabidopsis* RAR genes include homologues of many DNA repair genes that are defective in different human diseases. For example, hereditary breast cancer and nonpolyposis colon cancer, *Xeroderma pigmentosum*

(*D-XPD*, *F-XPF*, *B-ERCC3*), *Ataxia telangiectasia* (*ATM*), and Cockayne’s syndrome (*CSA*, *CSB*) (The *Arabidopsis* Genome Initiative, 2000). The presence of additional homologues of many RAR genes was found in *Arabidopsis* genome. This is seen for almost every major class of DNA repair, including recombination (four *RecA*), DNA ligation (four *DNA ligase I*), PR (five class I photolyase and one class II photolyase), and NER (six *RPA1*, two *RPA2*, two *Rad25*, three *TFB1*, and four *Rad23*). Regarding BER, *Arabidopsis* contained 16 homologues of DNA base glycosylates, which is the maximum number known for any species. It also contains three homologues of AP endonuclease. Some of the extra copies of RAR genes originated through relatively recent gene duplication because many of the sets of genes were more closely related to each other than to their homologues in any other species. The presence of duplicate (paralogous) genes might be responsible for greater repair capacity.

The multiplicity of RAR genes could also be partly due to the transfer of genes from the organellar genomes to the nucleus. Repair gene homologues that appear to be of plastid origin include the recombination protein *RecA*, *RecG*, and *SMS*, two class I photolyase homologues, *Fpg*, two *MutS2* proteins, and the transcription repair coupling factor *Mfd*. The homologues of *E. coli* repair gene *UvrABC* was found to be absent in *Arabidopsis*. As *Arabidopsis* showed many similarities to the set of DNA repair genes found in other eukaryotes, it will not be difficult to determine the functions of these gene-encoded proteins in the plant system (The *Arabidopsis* Genome Initiative, 2000). Recently, the *A. thaliana Rad50* homologue (*AtRad50*) has been isolated and shown to be lo-

cated on chromosome 2 (Gallego et al., 2001). The Rad50 protein is involved in the cellular response to DNA double-strand break (DSBs), including the detection of damage, activation of cell-cycle checkpoints, and DSB repair via recombination. Gallego et al. (2001) have shown that Rad50 homologue protein is involved in meiosis and DNA repair in plants.

VII. CONCLUSIONS AND FUTURE PROSPECTS

The survival of organisms depends on the accurate transmission of genetic information from one cell to its daughters. Such faithful transmission requires not only extreme accuracy in the replication of DNA and the precision in chromosome distribution, but also the ability to survive spontaneous and induced DNA damage while minimizing the number of heritable mutations. To achieve this fidelity, cells have evolved several DNA repair mechanisms that can monitor the structure of chromosome and coordinate repair and cell-cycle progression. These repair mechanisms also help in maintaining the genome integrity and in regulating mutation rates. If the DNA damage is not repaired properly, it can lead to various complications, including cell death or mutations. The mutagenicity may be caused by persistent, unrepaired adducts that are subsequently converted to mutations through DNA replication or to mutation fixation through misrepair.

In general, the damaged proteins and RNA molecules can be replaced using information encoded in the DNA, but the DNA molecules themselves are irreplaceable. Therefore, the damaged DNA

has to be repaired. DNA repair is possible largely because the DNA molecule consists of two complementary strands. DNA damage in one strand can be removed and accurately replaced by using the undamaged complementary strand as a template. DNA can become damaged by a variety of means, some spontaneous, others catalyzed by environmental agents or defect in replication, which can leave mispaired bases. The chemistry of DNA damage is diverse and complicated. Consequently, the cellular response to this damage includes a wide range of enzymatic systems that catalyze some of the most interesting chemical transformations in DNA metabolism.

During the last several years, considerable progress has been made in elucidating the components of the eukaryotic DNA damage response and DNA repair pathways. How damaged DNA activated DNA damage response proteins is a question of central importance to be addressed in near future. In principle, a sensor protein should have the ability to interact with damaged DNA. As there are many ways to damage DNA, it is important to determine whether there is a sensor for each type of damage. If the damage itself is sensed through the interaction with a sensor protein, this sensor protein may potentially interfere with repair proteins that also need access to damaged DNA. The cell must be aware not only of damage, but also of when the damage is repaired, because the completion of repair should be the signal for termination of the damage response. Further work is needed to define the activation of damage-signaling kinases. Another area ripe for future exploration is the relationship between the DNA damage signaling pathway and telomeric DNA structures. Any DNA

lesions that block the movement of DNA and RNA polymerases may kill cells by interfering with replication in actively growing and dividing cells or by blocking transcription and thus depriving non-dividing cells of an essential protein. Current models dealing with replication of damaged DNA postulate the initial dissociation of replicative DNA polymerases when encountering sites of DNA damage. Replication can then restart beyond lesions, or DNA polymerases specialized in translesion synthesis can temporarily replace replicative polymerases to synthesize across the lesion; replicative enzymes then take over again. The DNA helicase can help in this bypass reaction; however, this helicase-mediated bypass hypothesis is still speculation.

In modern biology the DNA repair study occupies a central place. The DNA repair field in eukaryotes has a strong connection with replication, transcription, recombination, and cell cycle control and checkpoints (G1/S and G2/M). Checkpoints are mechanisms that allow cells to deal with DNA damage or other insults. A major role of checkpoints is to delay cell cycle transitions in order to allow time for the damage to be repaired. Therefore, it is important to understand how checkpoints regulate the basic cell cycle machinery. The overall progress of research on DNA repair in plant cell has been very slow when compared with other systems. However, now this field has started progressing fast in plants especially after the isolation of several repair protein and genes. However, the twentieth century began with the rediscovery of Mendel's rules of inheritance in pea (De Vries, 1900), and it ends with the elucidation of the complete sequence of *Arabidopsis*. The analysis of completed sequence provides insights into

the genetic basis of the similarities and differences of diverse multicellular organisms. Regarding DNA repair genes, it appears that they are highly conserved between plants and mammals than within the animal kingdom, perhaps reflecting common factors such as DNA methylation.

DSB repair in plants is error prone, with the main pathway of repair being illegitimate recombination. Such an error-prone DNA repair mechanism might have an important impact on plant evolution. In plant the DSB repair products have largely been characterized as excision products of transposable elements or insertion products of T-DNAs. These products are similar to mammalian DSB repair products, which also suggests that plant DSB repair is similar to mammals. There is an urgent need to study the Ku and DNA-PK in plants in order to understand the mechanism of DSB repair. Still there is a need to identify all the repair genes in plants as reported in other systems. One way of identifying the genes involved in DNA repair is to study mutants that are sensitive (or resistant) to DNA-damaging agents. The identification of mutants that are selectively altered in only one type of DSB repair will shed light on the interrelations between the different DSB repair pathways. It might also allow the selective activation of a homologous pathway to achieve gene targeting.

There is still much to be done. Various models proposed for repairs are largely hypothetical and require further experimental support. Cloning of all the repair genes and studying knockouts or overexpression of these genes could provide a solution for a longstanding problem of gene targeting in plants. Further, the crystallographic studies of DNA repair enzymes will reveal a wide range of

structural motifs used to recognize specific lesions in DNA. These studies will also help to unravel the catalytic mechanism by which these lesions are repaired. We hope that in future new technologies, such as gene machines for reverse genetics, sequence analysis of whole plant genomes, and utilization of DNA chips, will give a boost to unravel the mystery behind DNA repair mechanisms. Also in the near future proteomics will provide a wealth of protein-protein interaction data, which will add to the knowledge of repair proteins function. The challenge for future research is to uncover the exact molecular mechanisms in plant DNA repair system through which these repair proteins repair the DNA damage in cooperation with other proteins. More research is needed in plants to answer the many questions that still surround the repair proteins and their many targets.

ACKNOWLEDGMENT

The writing of this review was initiated during the visit of the senior author (NT) to the University of Melbourne under the Australia India Council program.

REFERENCES

- Aboussekhra, A., Biggerstaff, M., Shivji, M. K. K., Vilpo, J. A., Moncollin, V., Podust, V. N., Protic, M., Hubscher, U., Egly, J.-M., and Wood, R. D. 1995. Mammalian DNA nucleotide excision repair reconstituted with purified components. *Cell* **80**:859–868.
- Aboussekhra, A. and Thoma, F. 1999. TATA-binding protein promotes the selective formation of UV-induced (6-4)-photoproducts and modulates DNA repair in the TATA box. *EMBO J.* **18**:433–443.
- Agrawal, A. and Kale, R. K. 2001. Radiation induced peroxidative damage: mechanism and significance. *Indian J. Exp. Biol.* **39**:291–309.
- Ahmad, M., Jarillo, J. A., Klimczak, L. J., Landry, L. G., Peng, T., Last, R. L., and Cashmore, A. R. 1997. An enzyme similar to animal type II photolyases mediates photoreactivation in *Arabidopsis*. *Plant Cell* **9**:199–207.
- Anderson, C. W. 1993. DNA damage and the DNA-activated protein kinase. *Trends Biochem. Sci.* **18**:433–437.
- Au, K. G., Welsh, K., and Modrich, P. 1992. Initiation of methyl-directed mismatch repair. *J. Biol. Chem.* **267**:12142–12148.
- Bailly, C., Benamar, A., Corbineau, F., and Cume, D. 1996. Changes in malondialdehyde content and in superoxide dismutase, catalase and glutathione reductase activities in sunflower seeds as related to deterioration during accelerated aging. *Physiologia Plantarum* **97**:104–110.
- Balajee, A. A. and Bohr, V. A. 2000. Genomic heterogeneity of nucleotide excision repair. *Gene* **250**:15–30.
- Barb, W. G., Baxendale, J. H., George, P., and Hargrave, K. R. 1957. Hydroxyl radical generation by Fenton reaction. *Trans. Faraday Soc.* **47**:462–470.
- Barber, A. A. and Berheim, F. 1967. Loss of fluidity and break down of cell membrane: functions and transmembrane ionic gradient by lipid peroxidation. *Adv. Gerontol. Res.* **2**:355–363.
- Barnes, D. E., Stamp, G., Rosewell, I., Denzel, A., and Lindahl, T. 1998.

- Targeted disruption of the gene encoding DNA ligase IV leads to lethality in embryonic mice. *Curr. Biol.* **8**:1395–1398.
- Batschauer, A. 1993. A plant gene for photolyase: An enzyme catalyzing the repair of UV-light induced DNA damage. *Plant J.* **4**:705–709.
- Beauchamp, C. O. and Fridovich, I. 1973. Isozymes of superoxide dismutase from wheat germ. *Biochem. Biophys. Acta.* **317**:50–56.
- Bennett, R. A. O., Wilson, D. M., Wong, D., and Demple, B. 1997. Interaction of human apurinic endonuclease and DNA polymerase beta in the base excision repair pathway. *Proc. Natl. Acad. Sci. USA*, **94**:7166–7169.
- Beranek, D. T. 1990. Distribution of methyl and ethyl adducts following alkylation with monofunctional alkylating agents. *Mutat. Res.* **231**:11–30.
- Bessho, T., Sancar, A., Thompson, L. H., and Thelen, M. P. 1997. Reconstitution of human excision nuclease with recombinant XPF-ERCC1 complex. *J. Biol. Chem.* **272**:3833–3837.
- Bethke, P. C. and Jones, R. L. 2001. Cell death of barley aleurone protoplasts is mediated by reactive oxygen species. *Plant J.* **25**:19–29.
- Bialkowski, K. and Olinski, R. 1999. Oxidative damage to plant DNA in relation to growth conditions. *Acta Biochim Pol.* **46**:43–49.
- Blunt, T., Finnie, N. J., Tacciolo, G. E., Smith, G. C. M., Demengeot, J., Gottlieb, T. M., Mizuta, R., Vrghe, A. J., Alt, F. W., Jeggo, P. A., and Jackson, S.P. 1995. Defective DNA-dependent protein kinase activity is linked to V(D)J recombination and DNA repair defects associated with the murine scid mutation. *Cell* **80**:813–823.
- Bormman, J. F. 1989. Target sites of UV-B radiation in photosynthesis of higher plants. *J. Photochem Photobiol. B* **4**:145–158.
- Bornman, J. F., Reuber, S., Chen, Y-P., and Weissenboeck, G. 1997. Ultraviolet radiation as stress factor and the role of protective pigments. In: *Plants and UV-B. Responses to Environmental Change* (Lumsden, P. J., Ed.). Cambridge: Cambridge University Press, pp. 157–168.
- Bornman, J. F. and Teramura, A. H. 1993. Effects of UV-B radiation on terrestrial plants. In: Young, A. R., Bjorn, L. O., Moan, J., and Nultsch, W. (Eds.) *Environmental UV Photobiology*. pp. 427–471. Plenum Press, New York.
- Boubriak, I., Kargiolaki, H., Lyne, L., and Osborne, D. J. 1997. The requirement for DNA repair in desiccation tolerance of germinating embryos. *Seed Science Research* **7**:97–105.
- Britt, A. B. 1996. DNA damage and repair in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **47**:75–100.
- Britt, A. B. 1999. Molecular genetics of DNA repair in higher plants. *TIPS* **4**:20–25.
- Britt, A. B., Chen, J.-J., Wykoff, D., and Mitchell, D. 1993. A UV-sensitive mutant of *Arabidopsis* defective in the repair of pyrimidine-pyrimidinone (6–4) dimers. *Science* **261**:1571–1574.
- Cannon, G. C., Hedrick, L. A., and Heinhorst, S. 1995. Repair mechanisms of UV-induced DNA damage in soybean chloroplasts. *Plant Mol. Biol.* **29**:1267–1277.
- Carpenter, C. L. and Cantley, L. C. 1996. Phosphoinositide kinases. *Curr. Opin. Cell Biol.* **2**:153–158.
- Cheah, K. S. E. and Osborne, D. J. 1978. DNA lesions occur with loss of viabil-

- ity in embryos of ageing rye seeds. *Nature* **272**:593–599.
- Chen, J.-J., Mitchell, D. L., and Britt, A. B. 1994. A light-dependent pathway for the elimination of UV-induced pyrimidine (6–4) pyrimidine photoproducts in *Arabidopsis*. *Plant Cell* **6**:1311–1317.
- Chen, L., Trujillo, K., Sung, P., and Tomkinson, A. E. 2000. Interactions of the DNA ligase IV-Xrcc4 complex with DNA ends and the DNA-dependent protein kinase. *J. Biol. Chem.* **275**:26196–26205.
- Chu, G. 1997. Double-strand break repair. *J. Biol. Chem.* **272**:24097–24100.
- Costanzo, V., Robertson, K., Ying, C. Y., Kim, E., Avvedimento, E., Gottesman, M., Grieco, D., and Gautier, J. 2000. Reconstitution of an ATP-dependent checkpoint that inhibits chromosomal DNA replication following DNA damage. *Molecular Cell* **6**:649–659.
- Coverley, D., Kenny, M. K., Lane, D. P., and Wood, R. D. 1992. A role for the human single-stranded DNA binding protein HSSB/RPA in an early stage of nucleotide excision repair. *Nucleic Acids Res.* **20**:3873–3880.
- Critchlow, S. E. and Jackson, S. P. 1998. DNA end-joining; from yeast to man. *Trends Biochem. Sci.* **23**:394–498.
- Culligan, K. M. and Hays, J. B. 1997. DNA mismatch repair in plants. An *Arabidopsis thaliana* gene that predicts a protein belonging to the MSH2 subfamily of eukaryotic mutS homologs. *Plant Physiol.* **115**:833–839.
- Dandoy, E., Schyns, R., Deltour, R., and Verly, W. G. 1987. Appearance and repair of apurine/apyrimidinic sites in DNA during early germination of *Zea mays*. *Mutation Res.* **181**:57–60.
- De Laat, W. L., Jaspers, G. J., and Hoeijmakers, J. H. J. 1999. Molecular mechanism of nucleotide excision repair. *Gene Dev.* **13**:768–785.
- Del Maestro, R. F., Bjork, J., and Arfors, K. E. 1981a. Increase in microvascular permeability induced by enzymatically generated free radicals. I. *In vivo* study. *Microvasc. Res.* **22**:239–254.
- Del Maestro, R. F., Bjork, J., and Arfors, K. E. 1981b. Increase in microvascular permeability induced by enzymatically generated free radicals. II. Role of superoxide anion radical, hydrogen peroxide, and hydroxyl radical. *Microvasc Res.* **22**:255–270.
- Demple, B. and Harrison, L., 1994. Repair of oxidative damage to DNA — enzymology and biology. *Annu. Rev. Biochem.* **63**:915–948.
- Dennis, C. and Surridge, C. 2000. *A. thaliana* genome. *Nature* **408**:791.
- De Vries, H. 1900. Sur la loi de disjonction des hybrides. *C. R. Acad. Sc. Paris.* **130**:845–847.
- Dianov, G. L., Prasad, R., Wilson, S. H., and Bohr, V. A. 1999. Role of DNA polymerase beta in the excision step of long patch mammalian base excision repair. *J. Biol. Chem.* **274**:13741–13743.
- Dizdaroglu, M. 1992. Measurement of radiation-induced damage to DNA at the molecular level. *Int. J. Radiat. Biol.* **61**:175–183.
- Doetsch, P. W. 1995. What's old is new: an alternative DNA excision repair pathway. *Trends Biochem. Sci.* **20**:384–386.
- Doetsch, P. W., McCray, W. H., Jr., and Valenzuela, M. R. 1989. Partial purification and characterisation of an endonuclease from spinach that cleaves ultraviolet light-damaged duplex DNA. *Biochim. Biophys. Acta* **1007**:309–317.

- Dormandy, T.L. 1980. Free radical reaction in biological systems. *Ann. Rev. Coll. Surg. Engl.* **62**:188–194.
- Doutriaux, M-P., Couteau, F., Bergounioux, C., and White, C. 1998. Isolation and characterisation of the RAD51 and DMC1 homologs from *Arabidopsis thaliana*. *Mol. Gen. Genet.* **257**:283–291.
- Drapkin, R., Reardon, J. T., Ansari, A., Huang, J. C., Zawal, L., Ahn, K., Sancar, A., and Reinberg, D. 1994. Dual role of TFIIH in DNA excision repair and in transcription by RNA polymerases II. *Nature* **368**:769–772.
- Drew, M. C., He, C.-J., and Morgan, P. W. 2000. Programmed cell death and aerenchyma formation in roots. *Trends Plant Sci.* **5**:123–127.
- Drummond, J. T., Li, G. M., Longley, M. J., and Modrich, P. 1995. Isolation of an hMSH2–p160 heterodimer that restores DNA mismatch repair to tumor cells. *Science* **268**:1909–1912.
- Eastwood, A. C. and McLennan, A. G. 1985. Repair replication in ultraviolet light-irradiated protoplasts of *Daucus carota*. *Biochem. Biophys. Acta* **826**:13–19.
- Ehtesham, N. Z., Tuan-Nghia, P., Gaikwad, A., Sopory, S., and Tuteja, N. 1999. Calnexin from *Pisum sativum*: molecular cloning of the cDNA and characterization of the encoded protein. *DNA Cell Biol.* **18**:865–874.
- Eisen, J. A. and Hanawalt, P. C. 1999. A phylogenomic study of DNA repair genes, proteins and processes. *Mutat. Res.* **435**:171–213.
- Elder, R. H., Dell'Aquila, A., Mezzina, M., Sarasin, A., and Osborne, D. J. 1987. DNA ligase in repair and replication in the embryos of rye, *Secale cereale*. *Mutat. Res.* **181**:61–71.
- Elder, R. H., and Osborne, D. J. 1993. Function of DNA synthesis and DNA repair in the survival of embryos during early germination and in dormancy. *Seed Sci. Res.* **3**:43–53.
- Essrs, J., van Steeg, H., de Wit, J., Swagemaekrs, S. M. A., Vermeij, M., Hoeijmker, J. H. J., and Kanaar, R. 2000. Homologous and non-homologous recombination differentially affect DNA damage repair in mice. *EMBO J.* **19**:1703–1710.
- Esterbauer, H. and Cheeseman, K. H. 1990a. Determination of aldehydic lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods Enzymol.* **186**:407–421.
- Esterbauer, H., Dieber-Rotheneder, M., Waeg, G., Puhl, H., and Tatzber, R. 1990b. Endogenous antioxidants and lipoprotein oxidation. *Biochem. Soc. Trans.* **18**:1059–1061.
- Evans, E., Moggs, J. G., Hwang, J. R., Egly, J.-M., and Wood, R. D. 1997. Mechanisms of open complex and dual incision formation by human nucleotide excision repair factors. *EMBO J.* **16**:6559–6573.
- Fee, J. A. 1982. In: *Oxidases and Related Redox Systems* (King, T.E., Hanson, T.S., and Morrison, M., Eds). Pergamon Press, Oxford, pp. 101–132.
- Fink, S. P., Reddy, G. R., and Marnett, L. J. 1997. Mutagenicity in *E. coli* of the major DNA adduct derived from the endogenous mutagen malondialdehyde. *Proc. Natl. Acad. Sci. USA* **94**:8652–8657.
- Fortini, P., Pascucci, B., Parlanti, E., Sobol, R. W., Wilson, S. H., and Dogliotti, E. 1998. Different DNA polymerases are involved in the short- and long-patch base excision repair in mammalian cells. *Biochemistry* **37**:3575–3580.
- Fridovich, I. 1976. Oxygen radicals, hydrogen peroxide and oxygen toxicity. In:

- Free Radical in Biology* (Pryor, W.A., Ed), 1, Academic Press, NY, pp. 239–277.
- Friedberg, E. C. 1985. DNA Repair. New York, W.H. Freeman.
- Friedberg, E. C., Walker, G. C., and Siede, W. 1995. In: *DNA Repair and Mutagenesis*. ASM Press, Washington DC.
- Friedberg, E. C. and Wood, R. D. 1996. DNA excision repair pathways. In: *DNA Replication in Eukaryotic Cells*. Cold Spring Harbor Laboratory Press, pp. 249–266.
- Frosina, G., Fortini, P., Rossi, O., Carrozzino, F., Raspaglio, G., Cox, L. S., Lane, D. P., Abbondandolo, A., and Dogliotti, E. 1996. Two pathways for base excision repair in mammalian cells. *J. Biol. Chem.* **271**:9573–9578.
- Gallego, F., Fleck, O., Li, A., Wyrzykowska, J., and Tinland, B. 2000. At RAD1, a plant homologue of human and yeast nucleotide excision repair endonucleases, is involved in dark repair of UV damages and recombination. *Plant J.* **21**:507–518.
- Gallego, M. E., Jeanneau, M., Granier, F., Bouchez, D., Bechtold, N., and White, C. I. 2001. Disruption of the *Arabidopsis Rad50* gene leads to plant sterility and MMS sensitivity. *Plant J.* **25**:31–41.
- Gary, R., Ludwig, D., Cornelius, H., MacInnes, M., and Park M. 1997. The DNA repair endonuclease XPG binds to proliferating cell nuclear antigen (PCNA) and shares sequence elements with the PCNA-binding regions of FEN-1 and cyclin-dependent kinase inhibitor P21. *J. Biol. Chem.* **272**:24522–24529.
- George, J. W., Ghate, S., Matson, S. W., and Besterman, J. M. 1992. Inhibition of DNA helicase II unwinding and ATPase activities by DNA-interacting ligands. *J. Biol. Chem.* **267**:10683–10689.
- Gerlach, V. L., Aravind, L., Gotway, G., Schultz, R. A., Koonin, E. V. and Friedberg, E. C. 1999. Human and mouse homologs of *E. coli* DinB (DNA polymerase IV), members of the UmuC/DinB superfamily. *Proc. Natl. Acad. Sci. USA* **21**:11922–11927.
- Gibbs, P. E., McGregor, W. G., Maer, V. M., Nisson, P., and Lawrence, C. W. 1998. A human homolog of the *Saccharomyces cerevisiae* REV3 gene, which encodes the catalytic subunit of DNA polymerase zeta. *Proc. Natl. Acad. Sci. USA* **94**:6876–6880.
- Gidrol, X., Serghini, H., Noubhani, A., Mocquot, B., and Pradet, P. 1989. Biochemical changes induced by accelerated aging in sunflower seeds. I. Lipid peroxidation and membrane damage. *Physiologia Plantarum* **76**:591–597.
- Goodman, B. A., McPhail, D. B., and Lnehan, D. J. 1986. Oxygen-induced free radical in wheat roots. *Free Rad. Res. Commun.* **2**:173–178.
- Goodman, M. E., Creighton, S., Bloom, L. B., and Petruska, J. 1993. Biochemical basis of DNA replication fidelity. *Crit. Rev. Biochem. Mol. Biol.* **28**:83–126.
- Gorbalenya, A. E. and Koonin, E. V. 1993. Helicases: amino-acid sequence comparisons and structure-function relationships. *Curr. Biol.* **3**:419–429.
- Gorbunova, V. and Levy, A. A. 1997. Non-homologous DNA end-joining in plant cells is associated with deletions and filler DNA insertions. *Nucleic Acids Res.* **25**:4650–4675.
- Gorbunova, V. and Levy, A. A. 1999. How plants make ends meet: DNA double-strand break repair. *Trends Plant Sci.* **4**:263–269.

- Gorman, M. A., Morera, S., Rothwell, D. G., de La Fortelle, E., Mol, C. D., Tainer, J. A., Hickson, I. D., and Freemont, P. S. 1997. The crystal structure of the human DNA repair endonuclease HAP1 suggests the recognition of extra-helical deoxyribose at DNA abasic sites. *EMBO J.* **16**:6548–6558.
- Grilley, M., Griffith, J., and Modrich, P. 1993. Bidirectional excision in methyl-directed mismatch repair. *J. Biol. Chem.* **268**:11830–11837.
- Grilley, M., Welsh, K. M., Su, S. S., and Modrich, P. 1989. Isolation and characterization of the *E. coli* mutL gene product. *J. Biol. Chem.* **264**:1000–1004.
- Gu, H., Marth, J. D., Orban, P. C., Mossmann, H., and Rajewsky, K. 1994. Deletion of a DNA polymerase beta gene segment in T cells using cell type-specific gene targeting. *Science* **265**:103–106.
- Halliwell, B. 1987. Oxidative damage, lipid peroxidation and antioxidant protection in chloroplasts. *Chem. Phys. Lipids* **44**:327–340.
- Halliwell, B. 1990. How to characterize a biological antioxidant. *Free Radic. Res. Commun.* **9**:1–32.
- Halliwell, B. and Gutteridge, J. M. C. 1984. Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem. J.* **219**:1–4.
- Halliwell, B. and Gutteridge, J. M. C. 1986. Oxygen-free radicals and iron in relation to biology and medicine: some problems and concepts. *Arch. Biochem. Biophys.* **246**:501–514.
- Hang, B., Signer, B., Mrgison, G. P., and Elder, R. H. 1997. Targeted deletion of alkylpurine-DNA-N-glycosylase in mice eliminates repair of 1,*N*⁶-ethenoadenine and hypoxanthine but not of 3,*N*⁴-ethenocytosine or 8-oxoguanine. *Proc. Natl. Acad. Sci. USA* **94**:12869–12874.
- Harber, F. and Weiss, J. 1934. The catalytic decomposition of hydrogen peroxide by iron salt. *Proc. R. Soc. Lond. A. Math. Phys. Sci.* **147**:337–351.
- Harlow, G. R., Jenkins, M. E., Pittalwala, T. S., and Mount, D. W. 1994. Isolation of *uvh1*, an *Arabidopsis* mutant hypersensitive to ultraviolet light and ionizing radiation. *Plant Cell* **6**:227–235.
- Harrington, J. J. and Lieber, M. R. (1994) Functional domains within Fen-1 and Rad2 define a family of structure-specific endonucleases; implications for nucleotide excision repair. *Genes Dev.* **8**:1344–1355.
- He, Z., Henricksen, L. A., Wold, M. S., and Ingles, C. J. 1995. RPA involvement in the damage-recognition and incision step of nucleotide excision repair. *Nature* **374**:566–569.
- Heath, R. L. and Taylor, G. E., Jr. 1997. Physiological processes and plant responses to ozone exposure. *Ecol. Stud.* **127**:317–368.
- Hemler, M. E. and Lands, W. E. 1980. Evidence for a peroxide-initiated free radical mechanism of prostaglandin biosynthesis. *J. Biol. Chem.* **255**:6253–6261.
- Hendry, G. A. F., Price, A. H., and Brocklebank, K. J. 1989. Role of iron in chlorophyll destruction in stressed plants. *Mol. Aspects Med.* **11**:132–133.
- Hidema, J., Kumagai, T., and Sutherland, B. M. 2000. UV radiation-sensitive norin 1 rice contains defective cyclobutane pyrimidine dimer photolyase. *Plant Cell*, **12**:1569–1578.
- Hidema, J., Kumagai, T., Sutherland, J. C., and Sutherland, B. M. 1997. Ultraviolet B-sensitive rice cultivar deficient

- in cyclobutyl pyrimidine dimer repair. *Plant Physiol.* **113**:39–44.
- Hodges, D. M., Andrews, C. J., Johnson, D. A., and Hamilton, R. I. 1997. Antioxidant enzyme and compound responses to chilling stress and their combining abilities in differentially sensitive maize hybrids. *Crop Sci.* **37**:857–863.
- Hoeijmakers, J. H. J. 1993a. Nucleotide excision repair I: from *E. coli* to yeast. *Trends Genet.* **9**:173–177.
- Hoeijmakers, J. H. J. 1993b. Nucleotide excision repair II. from yeast to mammals. *Trends Genet.* **9**: 211–27.
- Hoeijmakers, J. H. J. and Bootsma, D. 1990. Molecular genetics of eukaryotic DNA excision repair. *Cancer Cells* **2**:311–320.
- Howland, G. P. 1975. Dark-repair of ultraviolet-induced pyrimidine dimers in the DNA of wild carrot protoplasts. *Nature* **254**:160–161.
- Huang, J. C., Svododa, D. L., Reardon, J. T., and Sancar, A. 1992. Human nucleotide excision nuclease removes thymine dimers from DNA by incising the 22nd phosphodiester bond 5' and the 6th phosphodiester bond 3' to the photodimer. *Proc. Natl. Acad. Sci. USA* **89**:3664–3668.
- Hubscher, U. and Thommes, P. 1992. DNA polymerase: in search of a function. *Trends Biochem. Sci.* **17**:55–58.
- Hunter, N., Chambers, S. R., Louis, E. J., and Borts, R. H. 1996. The mismatch repair system contributes to meiotic sterility in an interspecific yeast hybrid. *EMBO J.* **15**:1726–1733.
- Ikenaga, M. and Mabuchi, T. 1996. Photo-reactivation of endosperm mutations in maize. *Radiat. Bot.* **6**:165–169.
- Illades-Aguilar, B. and Setlow, P. 1994. Autoprocessing of the protease that degrades small, acid-soluble proteins of spores of *Bacillus* species is triggered by low pH, dehydration, and dipicolinic acid. *J. Bacteriol.* **176**:7032–7037.
- Inze, D. and Van Montagu, M. 1995. Oxidative stress in plants. *Curr. Opin. Biotechnol.* **6**:153–158.
- Jabs, T. 1999. Reactive oxygen intermediates as mediators of programmed cell death in plants and animals. *Biochem. Pharm.* **57**:231–245.
- Jackson, J. F. 1987. DNA repair in pollen. A review. *Mutat. Res.* **181**:17–29.
- Jackson, J. F. and Linskens, H. F. 1978. Evidence for DNA repair after ultraviolet irradiation of *Petunia hybrida* pollen. *Mol. Gen. Genet.* **161**:117–120.
- Jansen, M. A., Gaba, V., and Greenberg, B. M. 1998. Higher plants and UV-B radiation: balancing damage, repair and acclimatation. *Trends Plant Sci.* **3**:131–135.
- Jeggo, A. P., Taccioli, G. E., and Jackson, S. P. 1995. Menage a trois: double strand break repair, V(D)J recombination and DNA-PK. *Bioessays* **17**:949–957.
- Jiang, C. Z., Yen, C.-N., Cronin, K., Mitchell, D., and Britt, A. B. 1997. UV- and gamma-radiation sensitive mutants of *Arabidopsis thaliana*. *Genetics* **147**:1401–1409.
- Jin, H., Cominelli, E., Bailey, P., Parr, A., Mehrtens, F., Jones, J., Tonelli, C., Weisshaar, B., and Martin, C. 2000. Transcriptional repression by AtMyB4 controls production of UV-protecting sunscreens in *Arabidopsis*. *EMBO J.* **19**:6150–6161.
- Johnson, K. A., Fink, S. P., and Marnett, L. J. 1997. Repair of propanodeoxyguanosine by nucleotide excision repair *in vivo* and *in vitro*. *J. Biol. Chem.* **272**:11434–11438.

- Johnson, R. E., Kondratick, C. M., Prakash, S., and Prakash, L. 1999a. hRAD30 mutations in the variant form of *Xeroderma pigmentosum*. *Science* **285**:263–265.
- Johnson, R. E., Kovvali, G. K., Prakash, L., and Prakash, S. 1996. Requirement of the yeast Msh3 and Msh6 genes for Msh2-dependent genomic stability. *J. Biol. Chem.* **271**:7285–7288.
- Johnson, R. E., Prakash, S., and Prakash, L. 1999b. Efficient bypass of a thymine-thymine dimer by yeast DNA polymerase, Pol η . *Science* **283**:1001–1004.
- Johnson, R. E., Washington, M. T., Haracska, L., Prakash, S., and Prakash, L. 2000a. Eukaryotic polymerases ι and ζ act sequentially to bypass DNA lesions. *Nature* **406**:1015–1019.
- Johnson, R. E., Washington, M. T., Prakash, S., and Prakash, L. 1999c. Bridging the gap: a family of novel DNA polymerases that replicate faulty DNA. *Proc. Natl. Acad. Sci. USA* **96**:12224–12226.
- Johnson, R. E., Washington, M. T., Prakash, S., and Prakash, L. 2000b. Fidelity of human DNA polymerase Π . *J. Biol. Chem.* **275**:7447–7450.
- Junop, M. S., Modesti, M., Guarne, A., Ghirlando, R., Gellert, M., and Yang, W. 2000. Crystal structure of the Xrcc4 DNA repair protein and implications for end joining. *EMBO J.* **19**:5962–5970.
- Kaminaka, H., Morita, S., Tokumoto, M., Masumura, T., and Tanaka, K. 1999. Differential gene expressions of rice superoxide dismutase isoforms to oxidative and environmental stresses. *Free Radic. Res.* **31** (Suppl.):S219–S225.
- Kanofsky, J. R. and Sima, P. 1991. Singlet oxygen production from the reactions of ozone with biological molecules. *J. Biol. Chem.* **266**:9039–9042.
- Kehrer, J. P. 2000. The Haber-Weiss reaction and mechanisms of toxicity. *Toxicology* **149**:43–50.
- Kim, C., Snyder, R. O., and Wold, M. S. 1992. Binding properties of replication protein-A from human and yeast cells. *Mol. Cell. Biol.* **12**:3050–3059.
- Kley, D., Kleinman, M., Sandermann, H., Jr., and Krupa, S. 1999. Photochemical oxidants: State of the science. *Environ. Pollut.* **100**:19–24.
- Klimyuk, V. and Jones, J. 1996. At DMC1, the *Arabidopsis* homologue of the yeast DMC1 gene: characterization, transposon-induced allelic variation and meiosis-specific expression of a pAtDMC1: GUS fusion. *Plant J.* **11**:1–14.
- Klungland, A. and Lindahl, T. 1997. Second pathway for completion of human DNA base excision-repair; reconstitution with purified proteins and requirement for DNase IV (FEN1). *EMBO J.* **16**:3341.
- Kobayashi, T., Hota, Y., and Tabata, S. 1993. Isolation and characterization of a yeast gene that is homologous with a meiosis-specific cDNA from a plant. *Mol. Gen. Genet.* **237**:225–232.
- Kolodner, D. and Marsischky, G. T. 1999. Eukaryotic DNA mismatch repair. *Curr. Opin. Genet. Dev.* **9**:89–96.
- Kolodner, R. 1996. Biochemistry and genetics of eukaryotic mis-match repair. *Genes Dev.* **10**:1433–1442.
- Krokan, H. E., Standal, R., and Slupphaug, G. 1997. DNA glycosylases in the base excision repair of DNA. *Biochem. J.* **325**:1–16.
- Kubota, Y., Nash, R. A., Klungland, A., Schar, P., Brnes, D. E., and Lindahl, T. 1996. Reconstitution of DNA base

- excision-repair with purified human proteins: interaction between DNA polymerase beta and the XRCC1 protein. *EMBO J.* **15**:6662–6670.
- Landry, L. G., Stapleton, A. E., Lim, J., Hofman, P., Hays, J., Walbot, V., and Last, R. 1997. An *Arabidopsis* photolyase mutant is hypersensitive to ultraviolet-B radiation. *Proc. Natl. Acad. Sci. USA* **94**:328–332.
- Lawrence, C. W. and Hinkle, D. C. 1996. DNA polymerase zeta and the control of DNA damage-induced mutagenesis in eukaryotes. *Cancer Surv.* **28**:21–31.
- Lehmann, A. R. 1998. Dual functions of DNA repair genes: molecular, cellular and clinical implications. *BioEssays* **20**:146–155.
- Leprince, O., Deltour, R., Thorpe, P. C., Atherton, N. M., and Hendry, G. A. 1990. The role of free radicals and radical processing systems in loss of desiccation tolerance in germinating maize. *New Phytol.* **116**:573–580.
- Leprince, O., Hendry, G. A., Atherton, N. M., and Walters-Vertucci, C. W. 1996. Free radicals and metabolism associated with the acquisition and loss of desiccation tolerance in developing seeds. *Biochem. Soc. Trans.* **24**:451–455.
- Levine, A., Tenhaken, R., Dixon, R., and Lamb C. 1994. H₂O₂ from the oxidative burst orchestrates the plant hypersensitive response. *Cell* **79**:583–93.
- Li, L., Elledge, S. J., Peterson, C. A., Bales, E. S., and Legerski, R. J. 1994. Specific association between the human DNA repair proteins XPA and ERCC1. *Proc. Natl. Acad. Sci. USA* **91**:5012–5016.
- Lieber, M. R. 1991. Site-specific recombination in the immune system. *FASEB J.* **5**:2934–2944.
- Lieber, M. R. 1997. The FEN-1 family of structure-specific nucleases in eukaryotic DNA replication, recombination and repair. *BioEssays* **19**:233–240.
- Lin, J. J., Phillips, A. M., Herst, J. E., and Sancar, A. 1992. Active site of (A)BC excinuclease. II. Binding, bending and catalysis mutants of UvrB reveal a direct role in 3' and indirect role in 5' incision. *J. Biol. Chem.* **267**:17693–17700.
- Lin, J. J. and Sancar, A. 1992. Active site of (A)BC excinuclease. I. Evidence for 5' incision by UvrC through a catalytic site involving Asp399, Asp438, Asp466 and His538 residues. *J. Biol. Chem.* **267**:17688–17692.
- Lin, W., Xin, H., Zhang, Y., Wu, X., and Wang, Z. 1999. The human REV1 gene codes for a DNA template-dependent dCMP transferase. *Nucl. Acids Res.* **27**:4468–4475.
- Lindahl, T. 1993. Instability and decay of the primary structure of DNA. *Nature* **362**:709–710.
- Lindahl, T. and Barnes, D. E. 1992. Mammalian DNA ligases. *Annu. Rev. Biochem.* **61**:251–281.
- Lindahl, T. and Wood, R. D. 1999. Quality control by DNA repair. *Science* **286**:1897–1905.
- Liu, L. and Wells, P. G. 1995. DNA oxidation as a potential molecular mechanism mediating drug-induced birth defects: phenytoin and structurally related teratogens initiate the formation of 8-hydroxy-2'-deoxyguanosine *in vitro* and *in vivo* in murine maternal hepatic and embryonic tissues. *Free Radic. Biol. Med.* **19**:639–648.
- Liu, Z., Hossain, G. S., Islas-Osuna, M. A., Mitchell, D. L., and Mount, D. W. 2000. Repair of UV damage in plants by nucleotide excision repair: *Arabidopsis*

- UVH1 DNA repair gene is a homolog of *Saccharomyces cerevisiae* RAD1. *Plant J.* **21**:519–528.
- Livingstone-Zatchej, M., Meier, A., Suter, B., and Thoma, F. 1997. RNA polymerase II transcription inhibits DNA repair by photolyase in the transcribed strand of active yeast genes. *Nucl. Acids Res.* **25**:3795–3800.
- Luxford, C., Morin, B., Dean, R. T., and Davies, M. J. 1999. Histone H1 and other protein and amino acid hydroperoxides can give rise to free radicals that oxidize DNA. *Biochem. J.* **344**:125–134.
- Marintchev, A., Mullen, M. A., Maciejewski, M. W., Pan, B., Gryk, M. R., and Mullen, G. P. 1999. Solution structure of the single-strand break repair protein XRCC1 N-terminal domain. *Nature Struct. Biol.* **6**:884–893.
- Marnett, L. J. 1999a Lipid peroxidation-DNA damage by malondialdehyde. *Mutat. Res.* **424**:83–95.
- Marnett, L. J. 1999b. Chemistry and biology of DNA damage by malondialdehyde. *IARC Sci. Publ.* **150**:17–27.
- Masutani, C., Kusumoto, R., Yamada, A., Dohmae, N., Yokoi, M., Yuasa, M., Araki, M., Iwai, S., Takio, K., and Hanaoka, F. 1999. The XPV (xeroderma pigmentosum variant) gene encodes human DNA polymerase η . *Nature* **399**:700–704.
- Masutani, C., Sugasawa, K., Yanagisawa, J., Sonoyama, T., Ui, M., Enomoto, T., Takio, K., Tanaka, K., van der Spek, P. J., Bootsman, D., Hoeijmakers, J. H. J., and Hanaoka, F. 1994. Purification and cloning of a nucleotide excision repair complex involving the *Xeroderma pigmentosum* group C protein and a human homolog of yeast Rad23. *EMBO J.* **13**:1831–1843.
- Matsuda, T., Saijo, M., Kuraoka, I., Kobayashi, T., Nahatssu, Y., Nagai, A., Enjoji, T., Masutani, C., Sugasawa, K., Hanaoka, F., Yasui, A., and Tanaka, K. 1995. DNA repair protein XPA binds to replication protein A (RPA). *J. Biol. Chem.* **270**:4152–4157.
- Matsumoto, Y. and Kim, K. 1995. Excision of deoxyribose phosphate residues by DNA polymerase beta during DNA repair. *Science* **269**:699–702.
- Matsumoto, Y., Kim, K., and Bogenhagen, D. F. 1994. Proliferating cell nuclear antigen-dependent a basic site repair in *Xenopus laevis* oocytes: an alternative pathway of base decision DNA repair. *Mol. Cell. Biol.* **14**:6187–6197.
- Matsunaga, T., Mu, D., Park, C. H., Reardon, J. T., and Sancar, A. 1995. Human DNA repair excision nuclease. Analysis of the roles of the subunits involved in dual incisions by using anti-XPG and anti-ERCC1 antibodies. *J. Biol. Chem.* **270**:20862–20869.
- Mattimore, V., and Battista, J. R. 1996. Radioresistance of *Deinococcus radiodurans*: functions necessary to survive ionizing radiation are also necessary to survive prolonged desiccation. *J. Bacteriol.* **178**:633–637.
- Mazur, S. J. and Grossman, L. 1991. Dimerization of *E. coli* UvrA and its binding to undamaged and ultraviolet light damaged DNA. *Biochemistry* **30**:4432–4443.
- McClennan, A. G. 1987. The repair of ultraviolet light-induced DNA damage in plant cells. *Mutat. Res.* **181**:1–7.
- McClintock, B. 1931. Cytological observations of deficiencies involving known genes, translocations and an inversion in *Zea mays*. *Mo. Agric. Exp. Res. Stn. Res. Bull.* **163**:4.
- McDonald, J. P., Rapic-Otrin, V., Epstein, J. A., Broughton, B. C., Wang, X.,

- Lehmann, A. R., Wolgemuth, D. J., and Woodgate, R. 1999. Novel human and mouse homologs of *Saccharomyces cerevisiae* DNA polymerase eta. *Genomics* **60**:20–30.
- Mehlhorn, H., Tabner, B., and Welburn, A. R. 1990. Electron spin resonance evidence for the formation of free radicals in plants exposed to ozone. *Phys. Plant* **79**:377–83.
- Mehlhorn, H. and Wellburn, A. R. 1987. Stress ethylene formation determines plant sensitivity to ozone. *Nature* **327**:417–418.
- Misra, H. P. and Fridovich, I. 1972. The role of superoxide anion in the autooxidation of epinephrin and a simple assay for superoxide dismutase. *J. Biol. Chem.* **247**:3170–3175.
- Modesti, M. Hesse, J. E., and Gellert, M. 1999. DNA binding of XRCC4 protein is associated with V(D)J recombination but not with stimulation of DNA ligase IV activity. *EMBO J.* **18**:2008–2018
- Modrich, P. 1994. Mismatch repair, genetic stability, and cancer. *Science* **266**:1959–1960.
- Moggs, J. G., Yarema, K. J., Essigmann, J. M., and Wood, R. D. 1996. Analysis of incision sites produced by human cells extracts and purified proteins during nucleotide excision repair of a 1, 3–intrastrand d(gpTpG)-C is platin adduct. *J. Biol. Chem.* **271**:7177–7186.
- Moore, M. H., Gulbis, J. M., Dodson, E. J., Demple, B., and Moody, P. C. E. 1994. Crystal structure of a suicidal DNA repair protein: the Ada *O*⁶-methylguanine-DNA methyltransferase from *E. coli*. *EMBO J.* **13**:1495–1501.
- Mu, D., Hsu, D.S., and Sancar, A. 1996. Reaction mechanism of human DNA repair excision nuclease. *J. Biol. Chem.* **271**:8285–8294.
- Mu, D., Park, C. H., Matsunaga, T., Hsu, D. S., Reardon, J. T., and Sancar, A. 1995. Reconstitution of human DNA repair excision nuclease in a highly defined system. *J. Biol. Chem.* **270**:2415–2418
- Mu, D. and Sancar, A. 1997. Model for XPC-independent transcription-coupled repair of pyrimidine dimers in humans. *J. Biol. Chem.* **272**:7570–7573.
- Mu, D., Turson, M., Duckett, D. R., Drummond, J. T., Modrich, P., and Sancar, A. 1997a. Recognition and repair of compound DNA lesions (base damage and mismatch) by human mismatch repair and excision repair systems. *Mol Cell Biol.* **17**:760–769.
- Mu, D., Wakasugi, M., Hsu, D. S., and Sancar, A. 1997b. Characterization of reaction intermediates of human excision repair nuclease. *J. Biol. Chem.* **272**:28971–28979.
- Mueller, J. P. and Smerdon, M. J. 1996. Rad23 is required for transcription-coupled repair and efficient overall repair in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **16**:2361–2368.
- Mustafa, M. G. and Tiernery, D.F. 1978. Biochemical metabolic changes in the lung and oxygen, ozone and nitrogen dioxide toxicity. *Ann. Rev. Resp. Dis.* **118**:1061–1090.
- Myles, G. M. and Sancar, A. 1991. Isolation and characterization of functional domains of uvrA. *Biochemistry* **30**:3834–3840.
- Nakajima, S., Sugiyama, M., Iwai, S., Hitorni, K., Otsoshi, E., Kim, S. T., Jiang, C. Z., Todo, T., Britt, A. B., and Yamamoto, K. 1998. Cloning and characterization of a gene (UVR3) required for photorepair of 6–4 products in *Arabidopsis thaliana*. *Nucl. Acids Res.* **26**:638–644.
- Naqvi, A., Chance, B., and Cadenos, E. 1986. Reactive oxygen intermediates in bio-

- chemistry. *Ann. Rev. Biochem.* **55**:137–166.
- Nelson, J. R., Lawrence, C. W., and Hinkle, D. C. 1996. Deoxycytidyl transferase activity of yeast REV1 protein. *Nature* **382**:729–731.
- Nuffer, M. G. 1957. Additional evidence on the effect of X-ray and ultraviolet radiation on mutation in maize. *Genetics* **42**:273–82.
- O'Donovan, A., Davies, A. A., Moggs, J. G., West, S. C., and Wood, R. D. 1994. XPG endonuclease makes the 3' incision in human DNA nucleotide excision repair. *Nature* **371**:432–435.
- Ohba, T., Yoshioka, Y., Machida, C., and Machida, Y. 1995. DNA rearrangement associated with the integration of T-DNA in tobacco: an example for multiple duplications of DNA around the integration target. *Plant J.* **7**:157–164.
- Orzaez, D. and Granell, A. 1997. DNA fragmentation is regulated by ethylene during carpel senescence in *Pisum sativum*. *Plant J.* **11**:137–144.
- Osborne, D. J. and Boubriak, I. 1994. DNA and desiccation tolerance. *Seed Science Res.* **4**:175–185.
- Overmyer, K., Tuominen, H., Kettunen, R., Betz, C., Langebartels, C., Sundermann, H., and Kangasjarvi, J. 2000. Ozone-sensitive *Arabidopsis rcd1* mutant reveals opposite roles for ethylene and jasmonate signaling pathways in regulating superoxide-dependent cell death. *Plant Cell* **12**:1849–1862.
- Painter, R. B. 1985. Inhibition and recovery of DNA synthesis in human cells after exposure to ultraviolet light. *Mutat. Res.* **145**:63–69.
- Palombo, F., Gallinari, P., Iaccarino, I., Lettieri, T., Hughes, M., DqArrigo, A., Truong, O., Hsuan, J. J., and Jiricny, J. 1995. GTBP, a 160–kilodalton protein essential for mismatch-binding activity in human cells. *Science* **268**:1912–1914.
- Palombo, F., Iaccarino, I., Nakajima, E., Ikejima, M., Shimada, T., and Jiricny, J. 1996. hMutS β , a heterodimer of hMSH2 and hMSH3, binds to insertion/deletion loops in DNA. *Curr. Biol.* **6**:1181–1184.
- Pang, Q. and Hays, J. B. 1991. UV-B inducible and temperature-sensitive photoreactivation of cyclobutane pyrimidine dimers in *Arabidopsis thaliana*. *Plant Physiol.* **95**:536–543.
- Parikh, S. S., Mol. C.D., Slupphaug, G., Bharati, S., Krokan, H. E., and Tainer, J. A. 1998. Base excision repair initiation revealed by crystal structures and binding kinetics of human uracil-DNA glycosylase with DNA. *EMBO J.* **17**:5214–5216.
- Park, C-H. and Sancar, A. 1994. Formation of a ternary complex by human XPA, ERCC1 and ERCC4 (XPF) excision repair proteins. *Proc. Natl. Acad. Sci. USA* **91**:5017–5021.
- Park, H. W., Kim, S. T., Sancar, A., and Deisenhofer, J. 1995. Crystal structure of DNA photolyase from *E. coli*. *Science* **268**:1866–1872.
- Pellinen, R., Palva, T., and Kangasjarvi, J. 1999. Subcellular localization of ozone-induced hydrogen peroxide production in birch (*Betula pendula*) leaf cells. *Plant J.* **20**:349–356.
- Petrini, J. H. J., Xiao, Y., and Weaver, D. T. 1995. DNA ligase I mediates essential functions in mammalian cells. *Mol. Cell Biol.* **15**:4303–4308.
- Pham, X. H., Reddy, M. K., Ehtesham, N. Z., Matta, B., and Tuteja, N. 2000. A DNA helicase from *Pisum sativum* is homologous to translation initiation

- factor and stimulates topoisomerase inactivity. *Plant J.* **24**:219–229.
- Pieper, R. O. DNA Repair in Higher eukaryotes. In: DNA Damage and Repair, vol. 2, J.A. Nickoloff and M. F. Hoekstra, Eds. (Humana Press, Totowa, NJ, 1998), pp. 33–49.
- Pinhero, R. G., Rao, M. V., Paliyath, G., Murr, D. P., and Fletcher, R. A. 1997. Changes in activities of antioxidant enzymes and their relationship to genetic and paclobutrizol-induced chilling tolerance of maize seedlings. *Plant Physiol.* **114**:695–704.
- Plooy, A. C., van Dijk, M., and Lohman, P. H. M. 1984. Induction and repair of DNA cross links in Chinese hamster ovary cells treated with various platinum coordination compounds in relation to platinum binding to DNA, cytotoxicity, mutagenicity and anti-tumor activity. *Cancer Res.* **44**:2043–2051.
- Podust, L. M., Podust, V. A., Floth, C., and Hubscher, U. 1994. Assembly of DNA polymerase δ and ϵ holoenzymes depends on the geometry of the DNA template. *Nucl. Acids Res.* **22**:2970–2975.
- Price, A. H., Atherton, N. M., and Hendry, G. A. F. 1989a. Plants under drought-stress generate activated oxygen. *Free Rad. Res. Commun.* **8**:61–66.
- Price, A. H., Atherton, N. M., and Hendry, G. A. F. 1989b. Superoxide formation and enhanced transition-metal uptake in drought-damaged wheat leaves. In: *Free Radicals, Diseased States and Antiradical Intervention* (Ed. by C. Rice-Evans) pp. 467–479, Richelieu Press, London.
- Price, A. H. and Hendry, G. A. F. 1989. Stress and the role of activated oxygen scavengers and protective enzymes in plants subjected to drought. *Biochem. Soc. Trans.* **17**:493–494.
- Priestley, D. A., Werner, B. G., Leopold, A. C., and McBride, M. G. 1985. Organic free radicals in seeds and pollen: the effect of hydration and aging. *Physiol. Plant.* **64**:88–94.
- Protic-Sabljić, M. and Kraemer, K. H. 1985. One pyrimidine dimer inactivates expression of a transfected gene in *Xeroderma pigmentosum* cells. *Proc. Natl. Acad. Sci. USA* **82**:6622–6626.
- Protic-Sabljić, M., Tuteja, N., Munson, P. J., Hauser, J., Kraemer, K. H., and Dixon, K. 1986. UV light-induced cyclobutane pyrimidine dimers are mutagenic in mammalian cells. *Mol. Cell. Biol.* **6**:3349–3356.
- Pryor, W.A. 1986. Oxy radical related species: their formation, life time and reactions. *Ann. Rev. Physiol.* **48**:657–667.
- Puchta, H. and Hohn, B. 1991. The mechanism of extrachromosomal homologous DNA recombination in plant cells. *Mol. Gen. Genet.* **230**:1–7.
- Quaite, F. E., Takayanagi, S., Ruffini, J., Sutherland, J. C., and Sutherland, B. M. 1994. DNA damage levels determine cyclobutyl pyrimidine dimer repair mechanisms in alfalfa seedlings. *Plant Cell* **6**:1635–1641.
- Rao, M. V. and Davis, K. R. 1999. Ozone-induced cell death occurs via two distinct mechanisms in *Arabidopsis*: the role of salicylic acid. *Plant J.* **17**:603–614.
- Reardon J. T., Mu D., and Sancar, S. 1996. Overproduction, purification and characterisation of the XPC subunit of the human DNA repair excision nuclease. *J. Biol. Chem.* **270**:2415–2418.
- Reuzeau, C. and Cavalie, G. 1995. Activities of free radical processing enzymes in dry sunflower seeds. *New Phytol.* **130**:59–66.

- Ribeiro, D. T., Machdo, C. R., Costa, R. M. A., Praekelt, U. M., Van-sluys, M. A. and Menck, C. F. M. 1998. Cloning of a cDNA from *Arabidopsis thaliana* homologous to the human XPB gene. *Gene* **208**: 207–213.
- Rinehart, T. A., Dean, C., and Weil, C. F. 1997. Comparative analysis of non-random DNA repair following *Ac* transposon excision in maize and *Arabidopsis*. *Plant J.* **12**:1419–1427.
- Rosen, D. R., Siddiqui, T., and Patterson, D. 1993. Mutation in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* **362**:59–62.
- Roy, R., Schaeffer, L., Humber, S., Vermeulen, W., Weeda, G., and Egly, J.-M. 1994. The DNA-dependent ATPase activity associated with the class II basic transcription factor BTF2/TFIIH. *J. Biol. Chem.* **269**:9862–9832.
- Saijo, M., Kuraoka, I., Masutani, C., Hanaoka, F., and Tanaka, K. 1996. Sequential binding of DNA repair proteins RPA and ERCC1 to XPA *in vitro*. *Nucl. Acids Res.* **24**:4719–4724.
- Sakumi, K. and Sekiguchi, M. 1990. Structures and functions of DNA glycosylases. *Mutat. Res.* **236**:161–62.
- Salomon, S. and Puchta, H. 1998. Capture of genomic and T-DNA sequences during double-strand break repair in somatic plant cells. *EMBO J.* **17**:6086–6095
- Sancar, A. 1994. Structure and function of DNA photolyase. *Biochemistry* **33**:2–9.
- Sancar, A. 1996. No ‘end of history’ for photolyases. *Science* **272**:48–49.
- Sancar, A. and Rupp, W. D. 1983. A novel repair enzyme: UVrABC excision nuclease of *Escherichia coli* cuts a DNA strand on both sides of the damaged region. *Cell* **33**:349–260.
- Sancar, G. B. 1990. DNA photolyases: physical properties, action mechanism and roles in dark repair. *Mutat. Res.* **236**:147–160.
- Sandermann, H., Jr., Ernst, D., Heller, W., and Langebartels, C. 1998. Ozone: an abiotic elicitor of plant defence reactions. *Trends Plant Sci.* **3**:47–50.
- Santerre, A. and Britt, A. B. 1994. Cloning of a 3-methyladenine-DNA glycosylase from *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **91**:2240–2244.
- Saparbaev, M. and Laval, J. 1998. 3,N⁴-ethenocytosine, a highly mutagenic adduct, is a primary substrate for *E. coli* double-stranded uracil-DNA glycosylase and human mismatch-specific thymine-DNA glycosylase. *Proc. Natl. Acad. Sci. USA* **95**:8508–8513.
- Satoh, M. S. and Lindahl, T. 1992. Role of poly(ADP-ribose) formation in DNA repair. *Nature* **356**:356–358.
- Satoh, M. S., Poirier, G. G., and Lindahl, T. 1993. NAD(+)-dependent repair of damaged DNA by human cell extracts. *J. Biol. Chem.* **268**:5480–5487.
- Schaeffer L., Moncollin, V., Roy, R., Staub, A., Mezzina, M., Sarasin, A., Weeda, G., Hoeijmakers, J. H. J., and Egly, J.-M. 1994. The ERCC2 DNA repair protein is associated with the class II BTF2/TFIIH transcription factor. *EMBO J.* **13**:2388–2392.
- Schaeffer, L., Roy, R., Humbert, S., Moncollin, V., Vermeulen, W., Hoeijmakers, J. H. J., Chambon, P., and Egly, J.-M. 1993. DNA repair helicase: a component of BTF2 (TFIIH) basic transcription factor. *Science* **260**:58–63.
- Schraudner, M., Moeder, W., Wiese, C., van Camp, W., Inze, D., Langebarts, C., and Sandermann, H., Jr. 1998. Ozone-induced oxidative burst in the ozone biomonitor plant, tobacco Bel W3. *Plant J.* **16**:235–245.

- Schultz, T. F. and Quatrano, R. S. 1997. Characterisation and expression of a rice RAD23 gene. *Plant Mol. Biol.* **34**:557–562.
- Scott, L., Lafoe, D., and Weil, C. F. 1996. Adjacent sequences influence DNA repair accompanying transposon excision in maize. *Genetics* **142**:237–246.
- Seeberg, E., Eide, L., and Bjoras, M. 1995. The base excision repair pathway. *Trends Biochem. Sci.* **20**:391–397.
- Selby, C. P. and Saner, P. 1993. Molecular mechanism of transcription-repair coupling. *Science* **260**:53–57.
- Senaratna, T. and McKersie, B. D. 1986. Loss of desiccation tolerance during seed germination: a free radical mechanism of injury. In: *Membranes, Metabolism and Dry Organisms* (Ed. by A. C. Leopold), pp. 85–102, Cornell University Press, Ithaca.
- Seroussi, E. and Lavi, S. 1993. Replication protein A is the major single-stranded DNA binding protein detected in mammalian cell extracts by gel retardation assays and UV cross-linking of long and short single-stranded DNA molecules. *J. Biol. Chem.* **268**:7147–7154.
- Setlow, B. and Setlow, P. 1996. Role of DNA repair in *Bacillus subtilis* spore resistance. *J. Bacteriol.* **178**:3486–3495.
- Setlow, P. 1992a. I will survive: protecting and repairing spore DNA. *J. Bacteriol.* **174**:2737–2741.
- Setlow, P. 1992b DNA in dormant spores of *Bacillus* species is in an A-like conformation. *Mol. Microbiol.* **6**:563–567.
- Sharma, Y. K. and Davis, K. R. 1997. The effects of ozone on anti-oxidant responses in plants. *Free Radical Biol. Med.* **23**:480–488.
- Shi, L., Kent, R., and Bence, A. B. 1997. Developmental expression of a DNA repair gene in *Arabidopsis*. *Mutat. Res.* **384**:145–146.
- Shiloh, Y. 1997. Ataxia-telangiectasia and the Nijmegen breakage syndrome: related disorders but genes apart. *Annu. Rev. Genet.* **31**:635–662.
- Shivji, M. K. K., Kenny, M. K., and Wood, R. D. 1992. Proliferating cell nuclear antigen is required for DNA excision repair. *Cell* **69**:367–374.
- Shivji, M. K. K., Podust, V. N., Hubscher, Y., and Wood, R. D. 1995. Nucleotide excision repair DNA synthesis by DNA polymerase epsilon in the presence of PCNA, RFC and RPA. *Biochemistry* **34**:5011–5017.
- Sies, H. 1991. *Oxidative Stress: Oxidant and Antioxidants*. Academic Press, London.
- Sobol, R. W., Horton, J. K., Kuhn, R., Gu, H., Singhal, R. K., Prasad, R., Rajewsky, K., and Wilson, S. H. 1996. Requirement of mammalian DNA polymerase-beta in base-excision repair. *Nature* **379**:183.
- Stapleton, A. E. 1992. Ultraviolet radiation and plants: burning questions. *Plant Cell* **4**:1353–1358.
- Stapleton, A. E., Thorner, C. S., and Walbot, V. 1997. UV-B component of sunlight causes measurable damage in field-grown maize (*Zea mays* L.): developmental and cellular heterogeneity of damage and repair. *Plant Cell Environ.* **20**:279–290.
- Sugasawa, K., Ng, J. M. Y., Masutani, C., van der Spek, P. J., Eker, A. P. M., Hanaoka, F., Bootsma, D., and Hoeijmakers, J. H. J. 1998. *Xeroderma pigmentosum* group C is the initiator of global genome nucleotide excision repair. *Mol. Cell* **2**:223–232.
- Sugawara, N. and Haber, J. E. 1992. Characterization of double-strand break-induced recombination: homology

- requirements and single-stranded DNA formation. *Mol. Cell Biol.* **12**:563–575.
- Sung, P., Bailly, V., Weber, C., Prakash, L., and Prakash, S. 1993. Human xeroderma pigmentosum group D gene encodes a DNA helicase. *Nature* **365**, 852–855.
- Suter, B., Livingstone-Zatchej, M., and Thoma, F. 1997. Chromatin structure modulates DNA repair by photolyase *in vivo*. *EMBO J.* **16**:2150–2160.
- Sutherland, B. M., Takayanagi, S., Sullivan, J. H., and Sutherland, J. C. 1996. Plant responses to changing environmental stress: cyclobutyl pyrimidine dimer repair in soybean leaves. *Photochem. Photobiol.* **64**:464–468.
- Svoboda, D. L., Taylor, J. S., Hearst, J. E., and Sancar, A. 1993. DNA repair by eukaryotic excision nuclease: removal of thymine dimer and psoralen monoadduct by HeLa cell free extract and of thymine dimer by *Xenopus laevis* oocytes. *J. Biol. Chem.* **268**:1931–1936.
- Swagemakers, S.M.A., Essers, J., Wit, J. De., Hoeijmakers, J.H.J., and Kanaar, R. 1998. The human Rad 54 recombinatorial DNA repair protein is a double-stranded DNA-dependent ATPase. *J. Biol. Chem.* **273**:28292–28297.
- Takata, M., Sasaki, M.S., Sonoda, E., Morrison, C., Hashimoto, M., Utsumi, H., Yamaguchi-Iwai, Y., Shinohara, A., and Takeda, S. 1998. Homologous recombination and non-homologous end-joining pathways of DNA double-strand break repair have overlapping roles in the maintenance of chromosomal integrity in vertebrate cells. *EMBO J.* **17**:5497–5508.
- Takeuchi, Y., Murakami, M., Nakaima, N., Kondo, N., and Nikaido, O. 1996. Induction and repair of damage to DNA in cucumber cotyledons irradiated with UVB. *Plant Cell Physiol.* **37**:181–187.
- Talpaert-Borle, M. and Liuzzi, M. 1982. Base-excision repair in carrot cells. *Eur. J. Biochem.* **124**:435–440.
- Tamada, T., Kitadokoro, K., Higuchi, Y., Inaka, K., Yasui, A., de Rooter, P. E., Eker, A. P., and Miki, K. 1997. Crystal structure of DNA photolyase from *Anacystis nidulans*. *Nature Struct. Biol.* **4**:887–891.
- Tang, M., Shen, X., Frank, E. G., O'Donnell, M., Woodgate, R., and Goodman, M. F. 1999. UmuD'(2)C is an error-prone DNA polymerase, *E. coli* pol V. *Proc. Natl. Acad. Sci. USA* **96**:8919–8924.
- Tappel, A. L. 1973. Lipid peroxidation damage to cell components. *Fed Proc.* **32**:1870–1874.
- Taylor, R.M. 1996. Ultraviolet B-induced DNA lesions and their removal in wheat (*Triticum aestivum* L.) leaves. *Plant Cell Environ.* **19**:171–181.
- Taylor, R. M., Nikaido, O., Jordan, B. R., Bray, C. M., and Tobin, A. K. 1996a. UV-B-induced DNA lesions and their removal in wheat (*Triticum aestivum* L.) leaves. *Plant Cell Environ.* **19**:171–181.
- Taylor, R. M., Rosamond, J., Tobin, A. K., and Bray, C. M. 1996b. The cloning and sequence analysis of DNA ligase from *Arabidopsis thaliana*. Accession Number:1359495.
- Teo, I., Sedgwick, B., Demple, B., Li, B., and Lindahl, T. 1984. Induction of resistance to alkylating agents in *E. coli*: the *ada+* ne product serves both as a regulatory protein and as an enzyme for repair of mutagenic damage. *EMBO J* **3**:2151–2157.
- Teramura, A. H. 1983. Effects of ultraviolet-B radiation on the growth and yield of crop plants. *Physiol. Plant* **58**:415–427.
- Teramura, A. H. and Sullivan J. H. 1994. Effects of UV-B radiation on photo-

- synthesis and growth of terrestrial plants. *Photosynth. Res.* **39**:463–473.
- Terasawa, M., Shinohara, A., Hotta, Y., Ogawa, H., and Ogawa, T. 1995. Localization of RecA-like recombination proteins on chromosomes of the lily at various meiotic stages. *Genes Dev.* **9**:925–934.
- Tevini, M. and Teramura, A. H. 1989. UV-B effects on terrestrial plants. *Photochem. Photobiol.* **50**:479–487.
- The Arabidopsis Genome Initiative. 2000. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**:796–815.
- Thoma, F. 1999. Light and dark in chromatin repair: repair of UV-induced DNA lesions by photolyase and nucleotide excision repair. *EMBO J.* **18**:6585–6598.
- Todo, T. 1999. Functional diversity of the DNA photolyase blue light receptor family. *Mutat. Res.* **434**:89–97.
- Trosko, J. E., and Mansour, V. H. 1969. Photoreactivation of ultraviolet light-induced pyrimidine dimers in Ginkgo cells grown *in vitro*. *Mutat. Res.* **7**:120–121.
- Turchi, J. J., Murante, R. S., and Bambara, R. A. 1992. DNA substrate specificity of DNA helicase E from calf thymus. *Nucleic Acids Res.* **20**:6075–6080.
- Tuteja, N. 1997. Unraveling DNA helicases from plant cells. *Plant Mol. Biol.* **33**:947–952.
- Tuteja, N. 2000. Plant Cell and viral helicases: essential enzymes for nucleic acid transactions. *Crit. Rev. Plant Sci.* **19**:449–478.
- Tuteja, N., Beven, A. F., Shaw, P. J., and Tuteja, R. 2001. A pea homologue of human DNA helicase I is localized within the dense fibrillar component of the nucleolus and stimulated by phosphorylation with CK2 and cdc2 protein kinases. *Plant J.* **25**:9–17.
- Tuteja, N., Ochem, A., Taneja, P., Tuteja, R., Skopac, D., and Falaschi, A. 1995. Purification properties of human DNA helicase VI. *Nucl. Acids Res.* **23**:2457–2463.
- Tuteja, N. and Phan, T. N. 1998. A chloroplast DNA helicase II from pea that prefers fork-like structures. *Plant Physiol.* **118**:1029–1039.
- Tuteja, N., Phan, T-N., and Tewari, K. K. 1996. Purification and characterization of a DNA helicase from pea chloroplast that translocates in the 3' to 5' direction. *Eur. J. Biochem.* **238**:54–63.
- Tuteja, N., Phan, T-N., Tuteja, R., Ochem, A., and Falaschi, A. 1997. Inhibition of DNA unwinding and ATPase activities of human DNA helicase II by chemotherapeutic agents. *Biochem. Biophys. Res. Commun.* **236**:636–640.
- Tuteja, N., Rahman, K., Tuteja, R., and Falaschi, A. 1991. DNA helicase IV from HeLa cells. *Nucl. Acids Res.* **19**:3613–3618.
- Tuteja, N., Rahman, K., Tuteja, R., and Falaschi, A. 1993. Human DNA helicase V, a novel DNA unwinding enzyme from HeLa cells. *Nucl. Acids Res.* **21**:2323–23329.
- Tuteja, N., Rahman, K., Tuteja, R., Ochem, A., Skopac, D., and Falaschi, A. 1992. DNA helicase III from HeLa cells: an enzyme that acts preferentially on partially unwound DNA duplexes. *Nucl. Acids Res.* **20**:5329–5337.
- Tuteja, N., and Tuteja, R. 1996. DNA helicases: the long unwinding road. *Nature Genetics* **13**:11–12.
- Tuteja, N. and Tuteja, R., 2001. Unraveling DNA repair in human: molecular mechanisms and consequences of

- repair defect. *Crit. Rev. Biochem. Mol. Biol.* (in press).
- Tuteja, N., Tuteja, R., Ochem, A., Taneja, P., Huang, N. W., Simoncsits, A., Susic, S., Rahman, R., Marusic, L., Chen, J., Zhang, J., Wang, S., Pongor, S., and Falaschi, A. 1994. Human DNA heliase II: a novel DNA unwinding enzyme identified as the Ku autoantigen. *EMBO J.* **13**:4991–5001.
- Tuteja, N., Tuteja, R., Rahman, K., Kang, L.-Y., and Falaschi, A. 1990. A DNA helicase from human cells. *Nucl. Acids Res.* **18**:6788–6792.
- Tuteja, R. and Tuteja, N. 2000. Ku autoantigen: a multifunctional DNA binding protein. *Crit. Rev. Biochem. Mol. Biol.* **35**:1–33.
- Urdea, M. S., Warner, B. D., Running, J. A., Stempien, M., Clyne, J., and Horn, T. 1988. A comparison of nonradioisotopic hybridization assay methods using fluorescen, chemiluminescent and enzyme labeled synthetic oligonucleotide probes. *Nucl. Acids Res.* **16**:4937–4955.
- Van Camp, W., Van Montagu, M., and Inze, D. 1994. Superoxide dismutases. In Causes of Photooxidative stress and amerlioration of defense systems in plants. Foyer, C. H. and Mullineaux, P. M., Eds., pp. 317–341. CRC Press, Boca Raton, FL.
- Vaughan, P., Lindahl, T., and Sedgwick, B. 1993. Induction of the adaptive response of *E. coli* to alkylation damage by the environmental mutagen, methyl chloride. *Mutat. Res.* **293**:249.
- Verhage, R. A., Zeeman, A. M., Lombaerts, M., van de Putte, P., and Brouwer, J. 1996. Analysis of gene- and strand-specific repair in the moderately UV-sensitive *Saccharomyces cerevisiae* rad23 mutant. *Mutat. Res.* **362**:155–165.
- Villani, G., and Tanguy Le Gac, N. 2000. Interactions of DNA helicases with damaged DNA: possible logical consequences. *J. Biol. Chem.* **275**:33185–33188.
- Vodenicharov, M. D., Sallmann, F. R., Satoh, M. S., and Poirier, G.G. 2000. Base excision repair is efficient in cells lacking poly(ADP-ribose) polymerase 1. *Nucl. Acids Res.* **28**:3887–3896.
- Vonarx, E. J., Mitchell, H. L., Karthikeyan, R., Chatterjee, I., and Kunz, B. A. 1998. DNA repair in higher plants. *Mutat. Res.* **400**:187–200.
- Wagner, J., Gruz, P., Kim, S.R., Yamada, M., Matsui, K., Fuchs, R.P., and Nohmi, T. 1999. The dinB gene encodes a novel *E. coli* DNA polymerase, DNA pol IV, involved in mutagenesis. *Mol. Cell.* **4**:281.
- Washington, M. T., Johnson, R. E., Prakash, S., and Prakash, L. 1999. Fidelity and processivity of *Saccharomyces cerevisiae* DNA polymerase η . *J. Biol. Chem.* **274**:36835–36838.
- Washington, M. T., Johnson, R. E., Prakash, S., and Prakash, L. 2000. Accuracy of thymine-thymine dimer bypass by *Saccharomyces cerevisiae* DNA polymerase η . *Proc. Natl. Acad. Sci. USA* **97**:3094–3099.
- Weisinger, R.A. and Fridovich, I. 1973. Mitochondrial superoxide dismutase site of synthesis and intermitochondrial localiztion. *J. Biol. Chem.* **248**:4793–4796.
- White, C. I. and Haber, J. E. 1990. Intermediates of recombination during mating type switching in *Saccharomyces cerevisiae*. *EMBO J* **9**:663–673.
- Wilson, D. M. and Thompson, L. H. 1997. Life without DNA repair. *Proc. Natl. Acad. Sci. USA* **94**:12754–12757.
- Winkler, G., Vermeulen, W, Coin, F., Egly, J., Hoeijmakers, J., and Weeda, G. 1998. Affinity purification of human

- DNA repair transcription factor TFIIH using epitope-tagged *Xeroderma pigmentosum* B protein. *J. Biol. Chem.* **273**:1092–1098.
- Wise, R. R. and Naylor, A. W. 1987. Chilling-enhanced photooxidation. The peroxidative destruction of lipids during chilling injury to photosynthesis and ultrastructure. *Plant Physiol.* **83**:272–277.
- Wood, R. D. 1997. Nucleotide excision repair in mammalian cells. *J. Biol. Chem.* **272**:23465–23468.
- Wood, R. D., Mitchell, M., Sgouros, J., and Lindahl, T. 2001. Human DNA repair genes. *Science*, **291**:1284–1289.
- Wu, X., Li, J., Li, X., Hsieh, C. L., Burgers, P. M., and Lieber, M. R. 1996. Processing of branched DNA intermediates by a complex of human FEN-1 and PCNA. *Nucl. Acids Res.* **24**:2036–2043.
- Wymann, M.P. and Pirola, L. 1998. Structure and function of phosphoinositide 3-kinases. *Biochim. Biophys. Acta* **1436**:127–150.
- Xu, H., Swobodo, I., Bhalla, P. L., Sijbers, A. M., Zhao, C., Ong, E.-K., Hoeijmakers, J. H. J., and Singh, M. B. 1998 Plant homologue of human excision repair gene ERCC1 points to conservation of DNA repair mechanisms. *Plant J.* **13**:823–829.
- Yang, H., Clendenin, W. M., Wong, D., Demple, B., Slupska, M. M., Chiang, J-H., and Miller, J. H. 2001. Enhanced activity of adenine-DNA glycosylase (Myh) by AP endonuclease (Ape1) in mammalian base excision repair of an A/GO mismatch. *Nucl. Acids Res.* **29**:743–752.
- Yasui, A. and Eker, A. P. M. 1998. DNA photolyases. In: *DNA Damage and Repair*, vol. II (Nickoloff, J. A. and Hoekstra, M. F., Eds.) Totowa: Humana Press, pp. 9–32.
- Yasui, A., Eker, A. P. M., Yasuhira, S., Yajima, H., Kobayashi, T., Takao, M., and Oikawa, A. 1994. A new class of DNA photolyases present in various organisms including aplacental mammals. *EMBO J.* **13**:6143–6151.
- Young, T. E., Gallie, D. R., and Demason, D. A. 1997. Ethylene-mediated programmed cell death during maize endosperm development of wild-type and *shrunk2* genotypes. *Plant Physiol.* **115**:737–751.
- Zhou, B.-B. and Elledge, S. J. 2000. The DNA damage response: putting checkpoints in perspective. *Nature* **408**:433–439.